



## **Cambridge International Examinations**

Cambridge International General Certificate of Secondary Education

| CANDIDATE<br>NAME |      |                     |                   |
|-------------------|------|---------------------|-------------------|
| CENTRE<br>NUMBER  |      | CANDIDATE<br>NUMBER |                   |
| CHEMISTRY         |      |                     | 0620/32           |
| Paper 3 (Extend   | ded) | Fe                  | bruary/March 2015 |
|                   |      |                     | 1 hour 15 minutes |

Candidates answer on the Question Paper.

No Additional Materials are required.

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



| I |     | ches the description.   |
|---|-----|---|
|   | (a) | an element which is gaseous at room temperature and pressure                                    |
|   |     | [1]   |
|   | (b) | an element that is added to water to kill bacteria  |
|   |     | [1]   |
|   | (c) | an element that forms a basic oxide of the type XO  |
|   | , D | [1]   |
|   | (a) | an element used as an inert atmosphere in lamps   |
|   | (e) | an element that forms an amphoteric oxide   |
|   | (0) | [1]   |
|   | (f) | an element that reacts vigorously with cold water to produce hydrogen                           |
|   |     | [1]   |
|   |     | [Total: 6]  |
| 2 | (a) | Define the term <i>isotope</i> .  |
|   |     |   |
|   |     | [2]   |
|   | (b) | The table gives information about four particles, <b>A</b> , <b>B</b> , <b>C</b> and <b>D</b> . |
|   | ` , | Complete the table.   |
|   |     | The first line has been done for you.   |
|   |     |   |

| particle | number of protons | number of electrons | number of neutrons | nucleon<br>number | symbol<br>or formula |
|----------|-------------------|---------------------|--------------------|-------------------|----------------------|
| Α        | 6                 | 6                   | 6                  | 12                | С                    |
| В        | 11                | 10                  | 12                 |                   |                      |
| С        | 8                 |                     | 8                  |                   | O <sup>2-</sup>      |
| D        |                   | 10                  |                    | 28                | Al <sup>3+</sup>     |

[7]

[Total: 9]

| 3 | Ammonia is manufactured by the Haber process. Nitrogen and hydrogen are passed over a catalyst |
|---|--|
|   | at a temperature of 450 °C and a pressure of 200 atmospheres.                                  |

The equation for the reaction is as follows.

$$N_2 + 3H_2 \rightleftharpoons 2NH_3$$

The forward reaction is exothermic.


| (a) | Sta   | te <b>one</b> use of ammonia.<br>[1   |
|-----|-------|---|
| (b) | Wh    | at is the meaning of the symbol <del>←</del> ?  |
| (c) | nitro | at are the sources of nitrogen and hydrogen used in the Haber process?  |
| (d) |       | rogen[2<br>ne the catalyst in the Haber process.  |
|     |       | [1  |
| (e) | (i)   | If a temperature higher than 450 °C was used in the Haber process, what would happen to the <b>rate</b> of the reaction? Give a reason for your answer. |
|     |       |   |
|     | (ii)  | If a temperature higher than 450 °C was used in the Haber process, what would happen to the <b>yield</b> of ammonia? Give a reason for your answer.     |
|     |       |   |
|     |       | ים  |

| (f) | (i)   | If a pressure higher than 200 atmospheres was used in the Haber process, what wou happen to the <b>yield</b> of ammonia? Give a reason for your answer. |     |
|-----|-------|---|-----|
|     |       |   |     |
|     | (ii)  | Explain why the rate of reaction would be faster if the pressure was greater the 200 atmospheres.   | an  |
|     |       |   |     |
|     | (iii) | Suggest <b>one</b> reason why a pressure higher than 200 atmospheres is not used in the Haber process.  | ιе  |
|     |       |   |     |
| (g) |       | aw a dot-and-cross diagram to show the arrangement of the outer (valency) electrons in or<br>lecule of ammonia.   | ıе  |
|     |       |   |     |
|     |       |   |     |
|     |       |   |     |
|     |       |   |     |
|     |       |   | [2] |
| (h) | Am    | monia acts as a base when it reacts with sulfuric acid.   |     |
|     | (i)   | What is a base?   | [1] |
|     | (ii)  | Write a balanced equation for the reaction between ammonia and sulfuric acid.   | 21  |
|     |       |   | .–, |

| (a) | A co | ompound <b>X</b> contains 82.76% of carbon by mass and 17.24% of hydrogen by mass. |     |
|-----|------|--|-----|
|     | (i)  | Calculate the empirical formula of compound <b>X</b> .                             |     |
|     |      |  |     |
|     | (ii) | Compound <b>X</b> has a relative molecular mass of 58.                             | [2] |
|     |      | Deduce the molecular formula of compound <b>X</b> .                                |     |
|     |      |  | [2] |
| (b) | Alke | enes are unsaturated hydrocarbons.   |     |
|     | (i)  | State the general formula of alkenes.  | F41 |
|     | (ii) | State the empirical formula of alkenes.  | [1] |
|     |      |  | [1] |
| (c) |      | at is meant by the term <i>unsaturated hydrocarbon</i> ?  aturated                 |     |
|     | hya  | rocarbon   |     |
|     |      |  | [2] |
|     |      |  |     |

| (d) | Describe a test that would distinguish between saturated and unsaturated hydrocarbons. |    |
|-----|--|----|
|     | reagent  |    |
|     | observation (saturated hydrocarbon)  |    |
|     | observation (unsaturated hydrocarbon)  |    |
|     |  | [3 |

(e) Addition polymers can be made from alkenes. The diagram shows part of an addition polymer.



| /i\ | Draw a circle on the diagram to show one ren | voot unit in this polymor | [4]     |
|-----|--|---------------------------|---------|
| (1) | Draw a circle on the diagram to show one rep | eat unit in this polymer. | _ [ ] ] |

(ii) Give the structure and the name of the monomer used to make this polymer. structure

| name | [2] |
|------|-----|


(iii) Give the structure of an isomer of the alkene in (e)(ii).

[1]

[Total: 15]


**5** Aluminium and iron are extracted from their ores by different methods.

Aluminium is extracted from its purified oxide ore by electrolysis.



| Πe  | aluminium   |     |
|-----|---|-----|
| (a) | What is the name of the ore of aluminium which consists mainly of aluminium oxide?  |     |
|     |   | [1] |
| (b) | The electrodes are both made of the same substance.   |     |
|     | Name this substance.  |     |
|     |   | [1] |
| (c) | Aluminium oxide is dissolved in molten cryolite before it is electrolysed.  |     |
|     | Give <b>two</b> reasons why aluminium oxide dissolved in molten cryolite is electrolysed rather the molten aluminium oxide alone. | ıan |
|     |   |     |
|     |   | [2] |
| (d) | Write the <b>ionic</b> equations for the reactions at the electrodes in this electrolysis.  |     |
|     | anode (positive electrode)  |     |
|     | cathode (negative electrode)  | [2] |
|     |   | 1   |

(e) Iron is extracted from its oxide ore by reduction using carbon in a blast furnace.



|     | (i)   | Place the elements aluminium, carbon and iron in order of reactivity with the <b>least</b> reactive element first. |
|-----|-------|--|
|     |       | [1]  |
|     | (ii)  | Use your answer to <b>(e)(i)</b> to explain why iron is extracted by reduction using carbon but aluminium is not.  |
|     |       |  |
|     |       | [1]  |
| (f) | Wh    | at is the name of the ore of iron which consists mainly of iron(III) oxide?  |
|     |       | [1]  |
| (g) | Wri   | te balanced equations for the reactions occurring in the blast furnace which involve                               |
|     | (i)   | the complete combustion of coke (carbon),  |
|     |       | [1]  |
|     | (ii)  | the production of carbon monoxide from carbon dioxide,   |
|     |       | [1]  |
|     | (iii) | the reduction of iron(III) oxide,  |
|     |       | [1]  |
|     | (iv)  | the formation of slag.   |
|     |       | [1]  |
|     |       |  |

**6** A student is told to produce the maximum amount of copper from a mixture of copper and copper(II) carbonate.

The student adds the mixture to an excess of dilute sulfuric acid in a beaker and stirs the mixture with a glass rod. The copper(II) carbonate reacts with the sulfuric acid, forming a solution of copper(II) sulfate but the copper does not react with the sulfuric acid.

## The student then

| • | removes | the unreacted | copper from | the mixture, |
|---|---------|---------------|-------------|--------------|
|---|---------|---------------|-------------|--------------|

| <ul><li>CC</li></ul> | onverts the | solution o | f copper( | II) s | sulfate | into | copper | bv a | series of | of reactions | 3. |
|----------------------|-------------|------------|-----------|-------|---------|------|--------|------|-----------|--------------|----|
|----------------------|-------------|------------|-----------|-------|---------|------|--------|------|-----------|--------------|----|

| (a) | sulf | scribe <b>two</b> things that the student would observe when the mixture is added to the dilute furic acid.   |
|-----|------|---|
|     |      | [2]   |
| (b) |      | scribe how the student can produce pure dry copper from the mixture of copper and $\operatorname{per}(\Pi)$ sulfate solution.   |
|     |      |   |
|     |      | [3]   |
| (c) |      | e student then adds sodium hydroxide solution to the copper(II) sulfate solution to produce $per(\mathrm{II})$ hydroxide.   |
|     | (i)  | Describe what the student would observe.  |
|     |      | [1]   |
|     | (ii) | Write an <b>ionic</b> equation for this reaction.   |
|     |      | [1]   |
| (d) |      | er separating the copper(II) hydroxide from the mixture, the copper(II) hydroxide is heated engly. The copper(II) hydroxide decomposes into copper(II) oxide and steam. |
|     | (i)  | Write an equation for the decomposition of $copper(II)$ hydroxide. Include state symbols.   |
|     |      | [2]   |
|     | (ii) | Name a non-metallic element that can be used to convert copper( $\mathrm{II}$ ) oxide into copper.  |
|     |      | [1]   |

© UCLES 2015 [Turn over

[Total: 10]

 $Ethanol\ is\ manufactured\ from\ glucose,\ C_6H_{12}O_6,\ by\ fermentation\ according\ to\ the\ following\ equation.$ 

|     |      | $C_6H_{12}O_6 \rightarrow 2C_2H_5OH + 2CO_2$   |     |
|-----|------|--|-----|
| (a) | Sta  | te the conditions required for this reaction.  |     |
|     |      |  |     |
|     |      |  | [2] |
| (b) | In a | an experiment, 30.0g of glucose was fermented.   |     |
|     | (i)  | Calculate the number of moles of glucose in 30.0 g.  |     |
|     |      |  |     |
|     |      |  |     |
|     |      | mol [  | [2] |
|     | (ii) | Calculate the maximum mass of ethanol that could be obtained from 30.0 g of glucose.   |     |
|     |      |  |     |
|     |      |  |     |
|     |      | g [  | [2] |
| (i  | iii) | Calculate the volume of carbon dioxide at room temperature and pressure that can lead to the control of the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that can lead to the carbon dioxide at room temperature and pressure that the carbon dioxide at the carbon dioxide a |     |
|     | •    | obtained from 30.0 g of glucose.   |     |
|     |      |  |     |
|     |      |  |     |
|     |      | dm³ [  | [1] |
| (c) | Eth  | anol can also be manufactured from ethene.   |     |
| . , | (i)  | Name the raw material which is the source of ethene.   |     |
|     | (')  |  | [1] |
|     | (ii) | Write a balanced equation for the manufacture of ethanol from ethene.  | 1   |
| ·   | - •  | ·<br>  | [1] |
|     |      | [Total:  | 9]  |

## **BLANK PAGE**

The Periodic Table of the Elements **DATA SHEET** 

|                       |   |   |                          |                             |                                   |                                   |                                  | 9.5                    | Group                              |                                      |                                   |                                      |                                    |                                    |                                   |                         |                                    |
|-----------------------|---|---|--------------------------|-----------------------------|-----------------------------------|-----------------------------------|----------------------------------|------------------------|------------------------------------|--------------------------------------|-----------------------------------|--------------------------------------|------------------------------------|------------------------------------|-----------------------------------|-------------------------|------------------------------------|
| _                     | =   |   |                          |                             |                                   |                                   |                                  |                        |                                    |                                      |                                   | =                                    | ≥                                  | >                                  | 5                                 | =>                      | 0                                  |
|                       |   |   |                          |                             |                                   |                                   | 1<br>Hydrogen                    |                        |                                    |                                      |                                   |                                      |                                    |                                    |                                   |                         | 4 <b>He</b> Helium                 |
| 7 Lithium 3 23        | Beryllium 4 24                                      |   |                          |                             |                                   |                                   |                                  |                        |                                    |                                      |                                   | 11<br>Boron<br>5                     | 12<br>Carbon<br>6                  | 14 <b>N</b> itrogen 7              | 16<br>Oxygen<br>8                 | 19 Fluorine 9 35.5      | Neon 10 40                         |
| Sodium 11             | Mg<br>Magnesium                                     |   |                          |                             |                                   |                                   |                                  |                        |                                    |                                      |                                   | At<br>Aluminium<br>13                | Silicon                            | Phosphorus                         | Sulfur<br>16                      | Chlorine                | Ar<br>Argon                        |
| 39 <b>K</b> Potassium | Calcium   | Scandium 21   | 48 <b>T</b> Titanium     | 51<br>Vanadium<br>23        | Chromium Chromium 24              | Mn<br>Manganese<br>25             | 56<br><b>Fe</b><br>Iron          | Cobalt                 | 59 <b>Ni</b> Nickel                | 64<br><b>Cu</b><br>Copper<br>29      | 65<br><b>Zn</b><br>30             | 70<br><b>Ga</b><br>Gallium<br>31     | 73<br><b>Ge</b><br>Germanium<br>32 | 75 <b>As</b> Arsenic 33            | Se Selenium 34                    | 80 <b>Br</b><br>Bromine | 84 <b>Kr</b> ypton 36              |
| Rb Rubidium           | St<br>Strontium                                     | 89 <b>Y</b>   | 2 <b>r</b> Zirconium 40  | Niobium Ni 41               | 96<br>Mo<br>Molybdenum<br>42      | Tc<br>Technetium<br>13            | Ru<br>Ruthenium<br>44            | Rhodium 45             | 106 Pd Palladium 46                | 108 <b>Ag</b> Silver 47              | Cadmium 48                        | 115 <b>In</b> Indium 49              | 119 <b>Sn</b> Tin 50               | 122<br><b>Sb</b><br>Antimony<br>51 | 128 <b>Te</b> Tellurium 52        | 127 <b>T</b> lodine 53  | 131<br><b>Xe</b><br>Xenon<br>54    |
| 133<br>Caesium<br>55  | 137 <b>Ba</b> Barium 56                             | 139 <b>La</b> Lanthanum *   | 178 <b>Hf</b> Hafnium 72 | 181 <b>Ta</b> Tanantalum    | 184<br><b>W</b><br>Tungsten<br>74 | 186<br><b>Re</b><br>Rhenium<br>75 | 190<br><b>Os</b><br>Osmium<br>76 | 192 <b>Ir</b> Iridium  | 195<br><b>Pt</b><br>Platinum<br>78 | 197<br><b>Au</b><br>Gold             | 201<br><b>Hg</b><br>Mercury<br>80 | 204 <b>T 1</b> Thallium 81           | 207 <b>Pb</b> Lead 82              | 209 <b>Bis</b> Bismuth 83          | Po<br>Polonium<br>84              | At<br>Astatine<br>85    | Radon<br>86                        |
| Francium<br>87        | 226<br><b>Ra</b><br>Radium<br>88                    | 227<br><b>Ac</b><br>Actinium †  | Ĺ                        |                             |                                   |                                   |                                  |                        |                                    |                                      |                                   |                                      |                                    |                                    |                                   |                         |                                    |
| *58-711<br>190-103    | *58-71 Lanthanoid series<br>190-103 Actinoid series | d series<br>series  |                          | 140 <b>Ce</b> Cerium 58     | Pr<br>Praseodymium<br>59          | Neodymium<br>60                   | Pm<br>Promethium<br>61           | Sm<br>Samarium<br>62   | 152<br><b>Eu</b><br>Europium<br>63 | 157<br><b>Gd</b><br>Gadolinium<br>64 | 159<br><b>Tb</b><br>Terbium<br>65 | 162<br><b>Dy</b><br>Dysprosium<br>66 | 165<br><b>Ho</b><br>Holmium<br>67  | 167<br><b>Er</b><br>Erbium<br>68   | 169<br><b>Tm</b><br>Thulium<br>69 | Yb<br>Ytterbium<br>70   | 175<br><b>Lu</b><br>Lutetium<br>71 |
| Key                   | a ×   | <ul> <li>a = relative atomic mass</li> <li>X = atomic symbol</li> <li>b = proton (atomic) number</li> </ul> |                          | 232<br><b>Th</b><br>Thorium | <b>Pa</b> Protactinium            | 238<br><b>U</b><br>Uranium<br>92  | Neptunium                        | <b>Pu</b> Plutonium 94 | Am<br>Americium<br>95              | <b>Cm</b><br>Curium                  | <b>BK</b><br>Berkelium<br>97      | Cf<br>Californium<br>98              | Es<br>Einsteinium<br>99            | Fm<br>Fermium<br>100               | Md<br>Mendelevium<br>101          | No<br>Nobelium<br>102   | <b>Lr</b><br>Lawrendum<br>103      |

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.