



## Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

			1 hour 15 minutes
Paper 4 Theory (Extended)		February/March	
CHEMISTRY			0620/42
CENTRE NUMBER		CANDIDATE NUMBER	
CANDIDATE NAME			

**READ THESE INSTRUCTIONS FIRST** 

No Additional Materials are required.

Candidates answer on the Question Paper.

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



- **1** This question is about gases.
  - (a) The following substances are gases at room temperature.


letter	Α	В	С	D	E	F	G	Н
substance	SO <sub>2</sub>	Ar	СО	Cl <sub>2</sub>	NH <sub>3</sub>	CO <sub>2</sub>	CH <sub>4</sub>	C <sub>3</sub> H <sub>8</sub>

Identify, by letter:

	(i)	a gas which combines with water to form acid rain	[1]
	(ii)	two gases which exist as diatomic molecules	[2]
	(iii)	a gas which bleaches damp litmus paper	[1]
	(iv)	a gas which is used as an inert atmosphere in lamps	[1]
	(v)	two gases which are found in clean dry air	[2]
	(vi)	two gases which are found in refinery gas.	[2]
(b)	NF <sub>3</sub>	has covalent bonds.	
	(i)	What is a covalent bond?	

(ii) Complete the dot-and-cross diagram to show the electron arrangement in a molecule of  ${\rm NF}_3$ .


Show outer shell electrons only.



[3]

(c) A	ir is a mixture. Nitrogen and oxygen are the two most common gases in air.
(	) What is meant by the term <i>mixture</i> ?
(i	) State the percentage of oxygen, to the nearest whole number, in clean dry air.
	[1]
(ii	) Describe the steps in the industrial process which enables nitrogen and oxygen to be separated from clean dry air.
	Use scientific terms in your answer.
	[3]
(iv	) Which physical property of nitrogen and oxygen allows them to be separated?
	[1]
	[Total: 20]

- 2 Sodium chloride is a typical ionic compound.
  - (a) The diagram shows part of a lattice of sodium chloride.
    - (i) Complete the diagram to show the ions present. Use '+' for Na $^+$  ions and '-' for C $l^-$  ions. One ion has been completed for you.



[2]

(ii)	How many electrons does a chloride ion have?	
		[1]
(iii)	Identify an element which has atoms with the same number of electrons as a sodium i	on.
		[1]
<b>(b)</b> Ele	ctrolysis of concentrated aqueous sodium chloride is an important industrial process.	
(i)	What is meant by the term <i>electrolysis</i> ?	
		[2]
(ii)	Name the products of the electrolysis of concentrated aqueous sodium chloride.	
	1	
	2	
	3	
(iii)	Write an ionic half-equation for the reaction at the cathode. Include state symbols.	[3]
		[2]

(c)	Silver chloride can be made by reacting aqueous sodium chloride with aqueous silver nitrate.
	The other product of the reaction is sodium nitrate. The chemical equation for the reaction is
	shown.

$$NaCl(aq) + AgNO_3(aq) \rightarrow AgCl(s) + NaNO_3(aq)$$

A student attempted to make the maximum amount of **sodium nitrate** crystals. The process involved three steps.

- **step 1** The student added aqueous sodium chloride to aqueous silver nitrate and stirred. Neither reagent was in excess.
- **step 2** The student filtered the mixture. The student then washed the residue and added the washings to the filtrate.
- **step 3** The student obtained sodium nitrate crystals from the filtrate.

(i)	Describe what the student observed in <b>step 1</b> .	
(ii)	Why was the residue washed in <b>step 2</b> ?	[1]
ii)	Give the names of the <b>two</b> processes which occurred in <b>step 3</b> .  1	
	2	

- (iv) The student started with 20 cm<sup>3</sup> of 0.20 mol/dm<sup>3</sup> NaCl(aq).
  - Determine the amount of NaCl(aq) used.

amount of NaCl(aq) used = ..... mol


The yield of NaNO<sub>3</sub> crystals was 90%.

Calculate the mass of NaNO<sub>3</sub> crystals made.

[Total: 21]

[2]

- **3** Limestone rock is mainly calcium carbonate, CaCO<sub>3</sub>.
  - (a) The 'limestone cycle' is shown. Each step is numbered.



(i)	Complete the box to give the chemical name and formula of lime.	[2]
(ii)	Which step involves a physical change?	
		[1]
(iii)	What type of reaction is <b>step 1</b> ?	
		[1]
(iv)	Suggest how step 2 could be reversed.	
		[1]
(v)	Write a chemical equation for <b>step 4</b> .	
		[1]
(vi)	Explain why <b>step 4</b> is a neutralisation reaction. Refer to the substances reacting in y answer.	our
		[2]

(b) Dolomite is a similar rock to limestone. Dolomite contains magnesium carbonate, MgCO<sub>3</sub>.

Write a chemical equation for the reaction between magnesium carbonate and dilute nitric acid.

(c)	Forsterite is another rock which contains a magnesium compound.
	A sample of forsterite has the following composition by mass: Mg, 2.73 g; Si, 1.58 g; O, 3.60 g.
	Calculate the empirical formula of forsterite.
	empirical formula =[2]

[Total: 12]

Amr	non	ia is an important chemical.	
(a) Ammonia is a base.			
	(i)	In chemistry, what is meant by the term base?	
	(ii)	Write a word equation to show ammonia behaving as a base.	[1]
		[	2
	Am	monia reacts with chlorine. The chemical equation is shown. $2NH_3(g) \ + \ 3Cl_2(g) \ \to \ N_2(g) \ + \ 6HCl(g)$ Calculate the volume of chlorine, measured at room temperature and pressure, needed react completely with 0.68 g of ammonia.	tc
		volume of chlorine =cm <sup>3</sup> [	3

	(	ii)	The chemica	l equation can	be represented	as shown
--	---	-----	-------------	----------------	----------------	----------

2 H—N—H + 3 C
$$l$$
—C $l$   $\rightarrow$  N $\Longrightarrow$ N + 6 H—C $l$ 

Use the bond energies in the table to determine the energy change,  $\Delta H$ , for the reaction between ammonia and chlorine.

bond	bond energy in kJ/mol
N–H	390
C1-C1	240
N≡N	945
H–C1	430

• energy needed to break bonds

	kJ
energy released when bonds are formed	

.....kJ

ullet energy change,  $\Delta H$ , for the reaction between ammonia and chlorine

		kJ [3]
(iii)	Is the reaction endothermic or exothermic? Explain your answer.	

(c) Ammonia reacts with oxygen at high temperatures in the presence of a suitable catalyst to form nitric oxide, NO.

	4	NH <sub>3</sub> (g) + 5C	$O_2(g) \rightarrow 4NO(g)$	$g) + 6H_2O(g)$		
(i)	Explain how this c	hemical equa	ation shows am	monia acting as	a reducing agen	t.
						[1]
(ii)	Suggest a suitable answer.	e catalyst for	the reaction fro	om the list of met	als. Give a reasc	n for your
	aluminium	calcium	platinum	potassium	sodium	
	suitable catalyst					
	reason					[21

[Total: 13]

5

Alc	ohol	s are a 'family' of organic molecules which have the same general formula.	
(a)		at is the name given to any 'family' of organic molecules which have the same gener nula and similar chemical properties?	al
		[	1]
(b)	Giv	e the general formula of alcohols.	
		[	1]
(c)	Pro	pan-1-ol can be made from propene.	
	(i)	Name the reagent and give the conditions needed to convert propene into propan-1-ol. reagent	
		conditions	 2]
	(ii)	Write a chemical equation for the complete combustion of propan-1-ol.	ر∠
		[	2]
(d)	As	imple sugar can be represented as shown.	
		н—о———о—н	
	Sim	ple sugars can be polymerised to make more complex carbohydrates.	
	(i)	Complete the diagram to show part of a carbohydrate <b>polymer</b> made from the simp sugar shown.	le
		]	2]
	(ii)	Name the chemical process which occurs when a carbohydrate polymer is broken dow into simple sugars.	/n
		[	1]
	(iii)	What conditions are needed for this process to occur?	
		[	1]

e)	Chr	omatography can be used to identify simple sugars in a mixture.
		tudent analysed a mixture of simple sugars by chromatography. All the simple sugars in the ture were colourless.
	(i)	What is the name given to the type of substance used to identify the positions of the simple sugars on the chromatogram?
		[1]
	(ii)	The student calculated the $R_{\rm f}$ value of a spot on the chromatogram.
		Complete the expression for the $R_{\rm f}$ value of the spot.
		$R_{\rm f}$ =
		[1]
(	(iii)	How could a student identify a simple sugar from its $R_{\rm f}$ value?
		[1]
(	(iv)	Sometimes not all the substances in a mixture can be identified from the chromatogram produced.
		Explain why this may happen.
		[1]

[Total: 14]

## **BLANK PAGE**

## **BLANK PAGE**

## **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The Periodic Table of Elements

110.7	<b>  </b>	۵ ۲	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	格	radon			
18.7	=			6	Щ	fluorine 19	17	Cl	chlorine 35.5	35	ğ	bromine 80	53	П	iodine 127	85	¥	astatine -			
3	>			00	0	oxygen 16	16	S	sulfur 32	34	Se	selenium 79	52	Тe	tellurium 128	84	Ъ	moloum –	116		livermorium -
;	>			7	z	nitrogen 14	15	۵	phosphorus 31	33	As	arsenic 75	51	Sb	antimony 122	83	<u>B</u>	bismuth 209			
2	≥			9	ပ	carbon 12	14	S	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Pb	lead 207	114	Εl	flerovium -
=	=			2	В	boron 11	13	Ρl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	11	thallium 204			
										30	Zu	zinc 65	48	р С	cadmium 112	80	Нg	mercury 201	112	S	copernicium -
										59	Cn	copper 64	47	Ag	silver 108	62	Au	gold 197	111	Rg	roentgenium -
eroup										28	Ż	nickel 59	46	Pd	palladium 106	78	చ	platinum 195	110	Ds	darmstadtium -
5				7						27	ပိ	cobalt 59	45	格	rhodium 103	77	ı	iridium 192	109	Ĭ	meitnerium -
		- I	hydrogen 1							26	Fe	iron 56	44	Ru	ruthenium 101	92	Os	osmium 190	108	Hs	hassium
							1			25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium
				_	loq	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
			Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	g	niobium 93	73	<u>a</u>	tantalum 181	105	В	
					atc	len 				22	j	titanium 48	40	Zr	zirconium 91	72	茔	hafnium 178	104	¥	rutherfordium -
										21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
:	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	56	Ba	barium 137	88	Ra	radium
	_			က	=	lithium 7	#	Na	sodium 23	19	×	potassium 39	37	В	rubidium 85	22	S	caesium 133	87	ᇁ	francium -

71	Γn	lutetium	175	103	ב	lawrencium	ı
70	Υp	ytterbium	173	102	9	nobelium	I
69	H	thulium	169	101	Md	mendelevium	I
89	ш	erbinm	167	100	Fm	ferminm	I
29	웃	holmium	165	66	Es	einsteinium	ı
99	۵	dysprosium	163	86	ర్	californium	ı
65	q	terbium	159	97	BK	berkelium	1
64	gq	gadolinium	157	96	Cm	curium	I
63	En	europium	152	92	Am	americium	ı
62	Sm	samarium	150	94	Pn	plutonium	I
61	Pm	promethium	ı	93	ď	neptunium	ı
09	βN	neodymium	144	92	$\supset$	uranium	238
69	Ā	praseodymium	141	91	Ра	protactinium	231
58	Ce	cerium	140	06	┖	thorium	232
22	Га	lanthanum	139	68	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).