

# **Cambridge IGCSE**<sup>™</sup>

| CANDIDATE<br>NAME |  |                     |  |  |
|-------------------|--|---------------------|--|--|
| CENTRE<br>NUMBER  |  | CANDIDATE<br>NUMBER |  |  |

CHEMISTRY 0620/32

Paper 3 Theory (Core)

February/March 2020

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

#### **INSTRUCTIONS**

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

#### **INFORMATION**

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].
- The Periodic Table is printed in the question paper.

1 (a) A list of compounds is shown.

aluminium oxide
calcium bromide
calcium oxide
ethane
ethene
hydrogen chloride
methane
nitrogen dioxide
potassium iodide
potassium manganate(VII)
sodium chloride


Answer the following questions using only the compounds in the list. Each compound may be used once, more than once or not at all.

## Which compound:

|            | (i)  | when in acidified solution, is used to test for sulfur dioxide                                    |     |
|------------|------|---|-----|
|            |      |   | [1] |
|            | (ii) | is the main constituent of natural gas  |     |
|            |      |   | [1] |
| <b>(</b> i | iii) | when dissolved in water, gives a yellow precipitate on addition of acidified aqued silver nitrate | ous |
|            |      |   | [1] |
| <b>(</b> i | iv)  | is used in flue gas desulfurisation to neutralise acidic gases                                    |     |
|            |      |   | [1] |
| (          | (v)  | is a reactant used in the manufacture of ethanol?   |     |
|            |      |   | [1] |
| (b)        | Wh   | at is the meaning of the term compound?   |     |
|            |      |   | [2] |

(c) Complete the electronic structure of a molecule of hydrogen chloride.

Show only the outer shell electrons.




[2]

[Total: 9]

2 Iron is extracted by heating a mixture of coke (carbon), limestone and iron ore in air in a blast furnace.

A diagram of the blast furnace is shown.



| (a) | Nar  | Name the ore of iron added to the blast furnace.                                      |         |  |  |  |  |  |
|-----|------|---|---------|--|--|--|--|--|
|     |      |   | [1]     |  |  |  |  |  |
| (b) | The  | e impurities in the iron ore are removed as slag.                                     |         |  |  |  |  |  |
|     | (i)  | What information in the diagram shows that slag is less dense than molten iron?       |         |  |  |  |  |  |
|     |      |   | [1]     |  |  |  |  |  |
|     | (ii) | Which one of the substances added to the blast furnace helps to remove the impurities | ∍s?     |  |  |  |  |  |
|     |      | Explain how it does this.   |         |  |  |  |  |  |
|     |      | substance   |         |  |  |  |  |  |
|     |      | explanation   |         |  |  |  |  |  |
|     |      |   |         |  |  |  |  |  |
|     |      |   | <br>[3] |  |  |  |  |  |
|     |      |   | [o]     |  |  |  |  |  |
| (c) | Hot  | air is blown into the blast furnace.  |         |  |  |  |  |  |
|     | Exp  | plain why.  |         |  |  |  |  |  |
|     |      |   |         |  |  |  |  |  |
|     |      |   | [1]     |  |  |  |  |  |

(d) The chemical equation for one of the reactions in the blast furnace is shown.

|     |      | $Fe_2O_3 + 3CO \rightarrow 2Fe + 3CO_2$  |            |
|-----|------|--|------------|
|     | (i)  | How does this equation show that Fe <sub>2</sub> O <sub>3</sub> has been reduced?  |            |
|     |      | [1   | ]          |
|     | (ii) | When 16.0 g of Fe <sub>2</sub> O <sub>3</sub> react with excess carbon monoxide, 11.2 g of iron are produced.  |            |
|     |      | Calculate the mass of iron produced when $4.0\mathrm{g}$ of $\mathrm{Fe_2O_3}$ react with excess carbon monoxide   | <b>)</b> . |
|     |      |  |            |
|     |      |  |            |
|     |      | mana of iron =   | 11         |
|     |      | mass of iron = g [1  | J          |
| (e) | An   | isotope of iron is shown.  |            |
|     |      | <sup>58</sup> Fe   |            |
|     | Dec  | duce the number of electrons, protons and neutrons in an atom of this isotope of iron.   |            |
|     | nun  | nber of electrons  |            |
|     | nun  | nber of protons  |            |
|     | nun  | nber of neutrons   |            |
|     |      |  | <i>i</i> ] |
| (f) | Iror | n is a transition element.   |            |
|     | Wh   | ich two of these statements about iron are correct?  |            |
|     | Tick | k two boxes.   |            |
|     |      | Iron forms coloured compounds.   |            |
|     |      | Iron can act as a catalyst.  |            |
|     |      | Iron is brown when freshly cut.  |            |
|     |      | Iron has a low density.  |            |
|     |      | Iron has a low melting point.  | >1         |
|     |      |  | _          |
| (f) | Wh   | ich <b>two</b> of these statements about iron are correct?  ( <b>two</b> boxes.  Iron forms coloured compounds.  Iron can act as a catalyst.  Iron is brown when freshly cut.  Iron has a low density. |            |
|     |      | [Total: 13   | 31         |

| Wa  | Water is essential for many industrial processes.                             |                  |  |  |  |  |  |  |
|-----|---|------------------|--|--|--|--|--|--|
| (a) | State <b>one</b> use of water in industry.                                    |                  |  |  |  |  |  |  |
|     |   | [1]              |  |  |  |  |  |  |
| (b) | What is the pH of pure water?   |                  |  |  |  |  |  |  |
|     | Draw a circle around the correct answer.                                      |                  |  |  |  |  |  |  |
|     | pH 0 pH 6 pH 7 pH 14  | [1]              |  |  |  |  |  |  |
| (c) | Filtration and chlorination are two of the steps used in water treatment.     |                  |  |  |  |  |  |  |
|     | Describe the purpose of each of these steps.                                  |                  |  |  |  |  |  |  |
|     | filtration  |                  |  |  |  |  |  |  |
|     |   |                  |  |  |  |  |  |  |
|     | chlorination  |                  |  |  |  |  |  |  |
|     |   |                  |  |  |  |  |  |  |
|     |   | [2]              |  |  |  |  |  |  |
| (d) | The changes of state of water are shown.                                      |                  |  |  |  |  |  |  |
|     | ice water steam   |                  |  |  |  |  |  |  |
|     | freezing  |                  |  |  |  |  |  |  |
|     | Give the names of the changes of state represented by <b>A</b> and <b>B</b> . |                  |  |  |  |  |  |  |
|     | A   |                  |  |  |  |  |  |  |
|     | В   | <br>[2]          |  |  |  |  |  |  |
|     |   | [ <del>-</del> ] |  |  |  |  |  |  |
|     |   |                  |  |  |  |  |  |  |

(e) The table compares the reactions of four metals with both steam and dilute hydrochloric acid.

| metal     | reaction with steam at 200 °C | observation with dilute hydrochloric acid |  |  |
|-----------|-------------------------------|---|--|--|
| copper    | no reaction                   | no bubbles formed bubbles form rapidly    |  |  |
| magnesium | rapid reaction                |   |  |  |
| nickel    | no reaction                   | bubbles form slowly                       |  |  |
| zinc      | rapid reaction                | bubbles form slowly                       |  |  |

Put the four metals in order of their reactivity. Put the least reactive metal first.

| least reactive — |  | most reactive |     |
|------------------|--|---------------|-----|
|                  |  |               |     |
|                  |  |               | [2] |

[Total: 8]

4 Angelic acid and ethanoic acid are both carboxylic acids.

The structure of angelic acid is shown.

- (a) (i) On the structure of angelic acid, draw a circle around the functional group that shows that it is a carboxylic acid. [1]
  - (ii) Deduce the formula of angelic acid to show the number of carbon, hydrogen and oxygen atoms.

| F A | -   |
|-----|-----|
| 11  |     |
|     |     |
|     | - 4 |

(iii) Angelic acid is an unsaturated compound.

Describe a chemical test to distinguish between an unsaturated and a saturated compound.

test .....

result with unsaturated compound .....

result with saturated compound ......[3]

(b) The formula of ethanoic acid is  $C_2H_4O_2$ .

Complete the table to calculate the relative molecular mass of ethanoic acid.

Use the Periodic Table to help you.

| type<br>of atom | number<br>of atoms | relative<br>atomic mass |             |
|-----------------|--------------------|-------------------------|-------------|
| carbon          | 2                  | 12                      | 2 × 12 = 24 |
| hydrogen        |                    |                         |             |
| oxygen          |                    | 16                      |             |

relative molecular mass = .....

[2]

(c) Ethanoic acid can be reduced to ethanol.

Complete the structure of ethanol to show all of the atoms and all of the bonds.

[1]

(d) Ethanol can be manufactured by fermentation.

| Des | scribe the process of fermentation to include:   |
|-----|--|
|     | the names of the reactants and catalyst  |
|     |  |
| •   | the conditions required  |
|     |  |
|     |  |
|     | the name of the process used to separate the ethanol from the rest of the reaction mixture |
|     |  |

[Total: 12]

[4]

5 The table shows some properties of four metals in Group I of the Periodic Table.

| metal     | melting point /°C | boiling point<br>/°C | relative electrical conductivity |
|-----------|-------------------|----------------------|----------------------------------|
| sodium    | 98                | 883                  |                                  |
| potassium | 63                | 760                  | 14                               |
| rubidium  |                   | 686                  | 8                                |
| caesium   | 29                | 669                  | 5                                |

| 1 | (a) | Com | olete | the | table | to | estima | te: |
|---|-----|-----|-------|-----|-------|----|--------|-----|
| ١ | a   |     | טוכנכ | uic | lanic | ιυ | Courna | ιc. |

| • | the | melting | point  | of | rubidiur |
|---|-----|---------|--------|----|----------|
| • | uic | mennig  | politi | Οı | rubiului |

• the relative electrical conductivity of sodium.

|     |      |   | [2] |
|-----|------|---|-----|
| (b) | Wh   | at is the physical state of caesium at 20°C?  |     |
|     | Giv  | e a reason for your answer.   |     |
|     |      |   |     |
|     |      |   |     |
|     |      |   | [2] |
| (c) | Des  | scribe the trend in the boiling points of the Group I metals.   |     |
|     |      |   | [1] |
| (d) |      | en potassium reacts with water, a coloured flame is seen and a gas is produced that pon a lighted splint. | ps  |
|     | (i)  | Complete the chemical equation for this reaction.   |     |
|     |      | K + $2H_2O \rightarrow 2KOH +$  | [2] |
|     | (ii) | State the colour of the flame when potassium reacts with water.   |     |
|     |      |   | [1] |
| (   | iii) | The solution formed is alkaline.  |     |
|     |      | Describe how you can use universal indicator solution to determine the pH of the soluti                   | on. |
|     |      |   |     |
|     |      |   |     |


[Total: 10]

| This   | question is about the re | eactions of magnes         | sium with nitric ac                                 | id.                       |       |
|--------|--------------------------|----------------------------|---|---------------------------|-------|
| (a) T  | he equation for the rea  | action of magnesiu         | m with concentra                                    | ted nitric acid is shown. |       |
|        | Mg                       | + 4HNO <sub>3</sub> → Mg(l | NO <sub>3</sub> ) <sub>2</sub> + 2NO <sub>2</sub> + | 2H <sub>2</sub> O         |       |
| (i     | ) The reaction is exc    | othermic.                  |   |                           |       |
|        | What is the meanir       | ng of the term exoti       | hermic?   |                           |       |
|        |                          |                            |   |                           | . [1] |
| (ii    | ) Which word best d      | escribes the compo         | ound Mg(NO <sub>3</sub> ) <sub>2</sub> ?            |                           |       |
|        | Draw a circle arour      | nd the correct answ        | ver.  |                           |       |
|        |                          | acid base                  | oxide s   | alt                       | [1]   |
| (iii   | ) Oxides of nitrogen     | are formed when for        | ossil fuels are bur                                 | rned.                     |       |
|        | What type of chem        | ical reaction occurs       | s when fossil fuels                                 | s are burned?             |       |
|        | Draw a circle arour      | nd the correct answ        | /er.  |                           |       |
|        | combustion               | cracking                   | fermentation  | neutralisation            | [1]   |
| (iv    | ) Oxides of nitrogen     | dissolve in rain wa        | ter to form acid ra                                 | ain.                      |       |
|        | State one adverse        | effect of acid rain        | on buildings.                                       |                           |       |
|        |                          |                            |   |                           | . [1] |
| (b) \/ | Vhen very dilute nitric  | acid reacts with ma        | agnesium nowder                                     | , hydrogen is produced.   |       |
|        | ·                        |                            |   |                           |       |
| (i     | ) Describe a practica    | ai memod for mves          | ligating the rate of                                | i tilis reaction.         |       |
|        |                          |                            |   |                           |       |
|        |                          |                            |   |                           |       |

| (ii) | What effect would each of the following have on the rate of this reaction? |    |
|------|--|----|
|      | Larger pieces of magnesium are used instead of magnesium powder.           |    |
|      | All other conditions stay the same.  |    |
|      |  |    |
|      | The temperature of the reaction mixture is increased.                      |    |
|      | All other conditions stay the same.  |    |
|      |  |    |
|      |  | [2 |

[Total: 9]

- 7 (a) The electrolysis of dilute sulfuric acid produces gases at both electrodes.
  - (i) The incomplete apparatus is shown.



Complete the diagram by:

- labelling the anode and cathode
- adding connecting wires
- showing how the gases are collected.


[3]

[2]


| (d) | Carbon dioxide is a greenhouse gas.                             |
|-----|---|
|     | State <b>one</b> effect of greenhouse gases on the environment. |
|     | [1]   |
|     | [Total: 10]   |

(a) The structures of four substances C, D, E and F, are shown. 8

C



Ε



- Which **one** of these substances, **C**, **D**, **E** or **F**, is a gas at room temperature? ......[1] What type of bonding is present in substance **E**? ......[1] (iii) Which one of these substances, C, D, E or F, is soluble in water? ......[1] (iv) Which one of these substances, C, D, E or F, conducts electricity when solid? .....[1]
- (b) The halogens have molecules containing two atoms. What is the name for molecules containing only two atoms?

| (c) | The  | e reaction of iodine with hydrogen is shown.  |      |
|-----|------|---|------|
|     |      | $I_2 + H_2 \rightleftharpoons 2HI$  |      |
|     | Wh   | at is the meaning of the symbol <del>←</del> ?                                      |      |
|     |      |   | [1]  |
|     |      |   |      |
| (d) | lod  | ine is formed when chlorine reacts with aqueous potassium iodide.                   |      |
|     | (i)  | Complete the chemical equation for this reaction.                                   |      |
|     |      | + 2KI $\rightarrow$ I <sub>2</sub> +KC $l$  | [2]  |
|     | (ii) | When aqueous iodine is mixed with aqueous potassium chloride, there is no reaction. |      |
|     |      | Suggest, in terms of chemical reactivity, why there is no reaction.                 |      |
|     |      |   | [1]  |
|     |      | [Total  | : 9] |

# **BLANK PAGE**

# **BLANK PAGE**

### **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

The Periodic Table of Elements

|       | = | 2<br>He | helium<br>4   | 10            | Ne           | neon<br>20                   | 18 | Ā          | argon<br>40      | 36 | 궃   | krypton<br>84   | 54 | Xe       | xenon<br>131     | 98    | 牊           | radon           |        |           |                    |
|-------|---|---------|---------------|---------------|--------------|------------------------------|----|------------|------------------|----|-----|-----------------|----|----------|------------------|-------|-------------|-----------------|--------|-----------|--------------------|
|       | = |         |               | 6             | ш            | fluorine<br>19               | 17 | Cl         | chlorine<br>35.5 | 35 | Ŗ   | bromine<br>80   | 53 | Н        | iodine<br>127    | 85    | ¥           | astatine<br>_   |        |           |                    |
|       | 5 |         |               | 80            | 0            | oxygen<br>16                 | 16 | ഗ          | sulfur<br>32     | 34 | Se  | selenium<br>79  | 52 | <u>a</u> | tellurium<br>128 | 84    | Ъ           | moloulum<br>—   | 116    |           | livermorium<br>–   |
|       | > |         |               | 7             | Z            | nitrogen<br>14               | 15 | ₾          | phosphorus<br>31 | 33 | As  | arsenic<br>75   | 51 | Sp       | antimony<br>122  | 83    | <u>.</u>    | bismuth<br>209  |        |           |                    |
|       | ≥ |         |               | 9             | ပ            | carbon<br>12                 | 14 | : <u>S</u> | silicon<br>28    | 32 | Эe  | germanium<br>73 | 20 | Sn       | tin<br>119       | 82    | Pb          | lead<br>207     | 114    | ŀΙ        | flerovium<br>-     |
|       | ≡ |         |               | 2             | М            | boron<br>11                  | 13 | Αl         | aluminium<br>27  | 31 | Ga  | gallium<br>70   | 49 | In       | indium<br>115    | 81    | lΤ          | thallium<br>204 |        |           |                    |
|       |   |         |               |               |              |                              |    |            |                  | 30 | Zn  | zinc<br>65      | 48 | g        | cadmium<br>112   | 80    | Ą           | mercury<br>201  | 112    | ű         | copernicium<br>-   |
|       |   |         |               |               |              |                              |    |            |                  | 29 | DO. | copper<br>64    | 47 | Ag       | silver<br>108    | 62    | Au          | gold<br>197     | 111    | Rg        | roentgenium<br>-   |
| Group |   |         |               |               |              |                              |    |            |                  | 28 | Z   | nickel<br>59    | 46 | Pd       | palladium<br>106 | 78    | చ           | platinum<br>195 | 110    | Ds        | darmstadtium<br>-  |
| ğ     |   |         |               | 1             |              |                              |    |            |                  | 27 | ပိ  | cobalt<br>59    | 45 | 格        | rhodium<br>103   | 77    | i           | iridium<br>192  | 109    | Ħ         | meitnerium<br>-    |
|       |   | - I     | hydrogen<br>1 |               |              |                              |    |            |                  | 26 | Pe  | iron<br>56      | 44 | Ru       | ruthenium<br>101 | 9/    | Os          | osmium<br>190   | 108    | Hs        | hassium            |
|       |   |         |               |               |              |                              | 1  |            |                  | 25 | M   | manganese<br>55 | 43 | ပ        | technetium<br>-  | 75    | Re          | rhenium<br>186  |        |           | bohrium<br>–       |
|       |   |         |               | _             | loq          | ass                          |    |            |                  | 24 | ပ်  | chromium<br>52  | 42 | Mo       | molybdenum<br>96 | 74    | ≥           | tungsten<br>184 | 106    | Sg        | seaborgium<br>-    |
|       |   |         | Key           | atomic number | atomic symbo | name<br>relative atomic mass |    |            |                  | 23 | >   | vanadium<br>51  | 41 | g        | niobium<br>93    | 73    | <u>a</u>    | tantalum<br>181 | 105    | Ор        | dubnium<br>—       |
|       |   |         |               |               | atc          | rel                          |    |            |                  | 22 | i   | titanium<br>48  | 40 | Zr       | zirconium<br>91  | 72    | Ξ           | hafnium<br>178  | 104    | Ŗ         | rutherfordium<br>- |
|       |   |         |               |               |              |                              |    |            |                  | 21 | Sc  | scandium<br>45  | 39 | >        | yttrium<br>89    | 57-71 | lanthanoids |                 | 89–103 | actinoids |                    |
|       | = |         |               | 4             | Be           | beryllium<br>9               | 12 | Mg         | magnesium<br>24  | 20 | Ca  | calcium<br>40   | 38 | ഗ്       | strontium<br>88  | 56    | Ba          | barium<br>137   | 88     | Ra        | radium             |
|       | _ |         |               | က             | =            | lithium<br>7                 | 1  | Na         | sodium<br>23     | 19 | ×   | potassium<br>39 | 37 | &<br>Q   | rubidium<br>85   | 55    | S           | caesium<br>133  | 87     | ъ         | francium<br>-      |

| 71<br>Lu        | lutetium<br>175     | 103 | ۲         | lawrencium   | ı   |
|-----------------|---------------------|-----|-----------|--------------|-----|
| 70<br>Yb        | ytterbium<br>173    | 102 | %         | nobelium     | ı   |
| °9<br>Tm        | thulium<br>169      | 101 | Md        | mendelevium  | ı   |
| 88<br>Fr        | erbium<br>167       | 100 | Fm        | ferminm      | ı   |
| 67<br>Ho        | holmium<br>165      | 66  | Es        | einsteinium  | ı   |
| °° O            | dysprosium<br>163   | 86  | ŭ         | californium  | ı   |
| e5<br>Tb        | terbium<br>159      | 97  | Ř         | berkelium    | ı   |
| Gd              | gadolinium<br>157   | 96  | Cm        | curium       | ı   |
| e3<br>Eu        | europium<br>152     | 92  | Am        | americium    | ı   |
| 62<br>Sm        | samarium<br>150     | 94  | Pu        | plutonium    | ı   |
| Pm              | promethium<br>-     | 93  | ď         | neptunium    | ı   |
| 9<br>9<br>N     | neodymium<br>144    | 92  | $\supset$ | uranium      | 238 |
| 59<br><b>Pr</b> | praseodymium<br>141 | 91  | Ра        | protactinium | 231 |
| Ce<br>Ce        | cerium<br>140       | 06  | 드         | thorium      | 232 |
| 57<br>La        | lanthanum<br>139    | 89  | Ac        | actinium     | ı   |

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).