



Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

Paper 4 Theory	(Extended)		May/June 2016
CHEMISTRY			0620/41
CENTRE NUMBER		CANDIDATE NUMBER	
CANDIDATE NAME			

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



1 hour 15 minutes

- 1 Protons, neutrons and electrons are subatomic particles.
 - (a) Complete the table to show the relative mass and relative charge of a proton, a neutron and an electron.

particle	relative mass	relative charge
proton		
neutron		
electron	<u>1</u> 1840	

b)	Bro	mine has two isotopes.	
	(i)	Define the term isotope.	
			[2
	(ii)	Explain why the two isotopes of bromine have the same chemical properties.	
			[2

(c) The table shows the number of protons, neutrons and electrons in some atoms and ions.
Complete the table.

particle	number of protons	number of neutrons	number of electrons
⁷ ₃Li			
³⁴ ₁₆ S ²⁻			
	19	22	18

[5]

[Total: 12]

- 2 Period 3 contains the elements sodium to argon. This question asks about the chemistry of each of the Period 3 elements or their compounds.
 - (a) Sodium nitrate is a white crystalline solid. When heated it melts and the following reaction occurs.

$$2NaNO_3(I) \rightarrow 2NaNO_2(I) + O_2(g)$$

A 3.40 g sample of sodium nitrate is heated.

Calculate the

number of moles of NaNO₃ used,

• number of moles of O₂ formed,

															mol

• volume of O₂ formed, in dm³ (measured at r.t.p.).

															(b	n	n	3
																	۲	3	1

- **(b)** Magnesium reacts slowly with warm water to form a base, magnesium hydroxide.
 - (i) Explain what is meant by the term base.

(ii) Write a chemical equation for the reaction between magnesium and warm water.




(c)	Alu	minium oxide is amphoteric. It is insoluble in water.
	Des	scribe experiments to show that aluminium oxide is amphoteric.
		[3]
(d)	Silid	con(IV) oxide has a giant structure.
	(i)	Name the type of bonding in silicon(IV) oxide.
	(ii)	Give two physical properties of silicon(IV) oxide.
		[2]
(e)		cium phosphate is used in fertilisers. The bonding in calcium phosphate is ionic. cium phosphate contains the phosphate ion, ${\rm PO_4}^{3-}$.
	(i)	What is ionic bonding?
		[2]
	(ii)	Deduce the formula of calcium phosphate.
		[1]

(f) Sulfur tetrafluoride, SF₄, can be made by combining gaseous sulfur with fluorine.

$$S(g) + 2F_2(g) \rightarrow SF_4(g)$$

The reaction is exothermic.

(i) Complete the energy level diagram for this reaction. Include an arrow which clearly shows the energy change during the reaction.



(ii) During the reaction the amount of energy given out is 780 kJ/mol.

The F–F bond energy is 160 kJ/mol.

Use this information to determine the bond energy, in kJ/mol, of one S-F bond in SF₄.

$$S + \begin{matrix} F - F \\ F - F \end{matrix} \rightarrow F - \begin{matrix} F \\ S - F \\ F \end{matrix}$$

..... kJ/mol [3]

[3]


(g)		orine and compounds of chlorine are important in water treatment and in laboratory testing water.
	(i)	Chlorine is added to water to make the water safe to drink.
		Explain why adding chlorine makes water safe to drink.
		[1]
	(ii)	A compound of chlorine is used in the laboratory to test for the presence of water.
		Name the compound of chlorine used in this test and describe the colour change seen in a positive result of this test.
		name of compound
		colour change from to
(h)	Arg	on is an unreactive noble gas.
	(i)	Explain why argon is unreactive.
		[1]
	(ii)	Give one use of argon.
		[1]
		[Total: 27]

Question 3 starts on the next page.

When aqueous sodium thiosulfate and dilute hydrochloric acid are mixed, a precipitate of insoluble sulfur is produced. This makes the mixture difficult to see through.

$$Na_2S_2O_3(aq) + 2HCl(aq) \rightarrow S(s) + 2NaCl(aq) + H_2O(l) + SO_2(g)$$

The time taken for the cross to disappear from view is measured.



A student adds the following volumes of aqueous sodium thiosulfate, dilute hydrochloric acid and distilled water to the conical flask.

The time taken for the formation of the precipitate of sulfur to make the cross disappear from view is recorded.


experiment number	volume of sodium thiosulfate /cm³	volume of hydrochloric acid /cm³	volume of distilled water /cm³	time taken for cross to disappear from view/s
1	10	10	40	56
2	20	10	30	28
3				

(a)	State the order in which the aqueous sodium thiosulfate, hydrochloric acid and distilled wate should be added to the flask.
	[1

(b) In experiment 3 the student wanted the sodium thiosulfate to be double the concentration used

	ın e	experiment 2.	
	(i)	Complete the table to show the volumes which should be used and the expected time taken for the cross to disappear from view in experiment 3.	e 2]
	(ii)	Use collision theory to explain why increasing the concentration of sodium thiosulfat would change the rate of reaction.	е
		[2	2]
(c)	The	e student repeated experiment 1 at a higher temperature.	
	Use	e collision theory to explain why the rate of reaction would increase.	
		[3]
		[Total: 8	3]

- **4** Electroplating steel objects with silver involves a three-step process.
 - **step 1** A coating of copper is applied to the object.
 - step 2 A coating of nickel is applied to the object.
 - **step 3** The coating of silver is applied to the object.
 - (a) A diagram of the apparatus used for **step 1** is shown.



(i) The chemical process taking place on the surface of the object is

Explain whether this process is oxidation or reduction.

$$Cu^{2+}(aq) + 2e^- \rightarrow Cu(s)$$

(ii)	Explain why the concentration of copper ions in the electrolyte remains constant throughout step 1 .

(b)	Give two changes which would be needed in order to coat nickel onto the object in step 2 .
	[2]
(c)	Copper, nickel and silver are transition elements. Typical physical properties of transition elements are a high density and a high melting point.
	Give three different properties of transition metals which are not typical of other metals.
	[3]
	[Total: 8]

Sul	furic acid is produced by the Contact process. The steps of the Contact process are shown.	
	starting material sulfur dioxide step 2 sulfur trioxide step 3 oleum step 4 sulfur acid	
(a)	Sulfur is a common starting material for the Contact process.	
	Name a source of sulfur.	
		[1]
(b)	Describe step 2 , giving reaction conditions and a chemical equation. Reference to react rate and yield is not required.	ion
		[5]
(c)	Step 3 involves adding sulfur trioxide to concentrated sulfuric acid to form oleum.	
	Complete the chemical equation for this reaction.	
	$H_2SO_4 + SO_3 \rightarrow \dots$	[1]

(d)	Dilu	te sulfuric acid is a typical acid.
		tudent adds excess dilute sulfuric acid to a sample of solid copper(II) carbonate in a -tube.
	(i)	Give three observations the student would make.
		[2]
	(ii)	Give the names of all products formed.
		[1]
(e)	Cor	ncentrated sulfuric acid has different properties to dilute sulfuric acid.
		en concentrated sulfuric acid is added to glucose, $C_6H_{12}O_6$, steam is given off and a black d is formed.
	(i)	Name the black solid.
		[1]
	(ii)	What type of reaction has occurred?
		[1]
		[Total: 12]

6

Petro	oleum is a source of many important chemicals.	
(a) I	Name two industrial processes which must take place to produce alkenes from petroleum.	
		[2]
(b) i	Ethene, CH ₂ =CH ₂ , and propene, CH ₂ =CHCH ₃ , can both be converted into polymers.	
((i) What type of polymerisation takes place when ethene forms a polymer?	
		[1]
(i	ii) What is the empirical formula of the polymer formed from ethene?	
		[1]
(ii	ii) Propene has the structural formula CH ₂ =CHCH ₃ .	
	Draw two repeat units of the polymer made from propene.	
		[2]
(c) l	Ethene will react with steam to form ethanol.	
ŀ	Propene will react with steam to form two isomers, both of which are alcohols.	
,	Suggest the structures of these alcohols.	

d)		ers are organic chemicals noted for their characteristic smells. Ethanoic acid and methanoreact to form an ester.	ol
	(i)	Name the catalyst needed to form an ester from ethanoic acid and methanol.	
		[1	1]
	(ii)	Name the ester formed when ethanoic acid reacts with methanol.	
		[1	1]
(iii)	Draw the structure of the ester formed when ethanoic acid reacts with methanol. Show a bonds.	ıII
		[2	2]
(iv)	Give the name of a polyester.	
		[1	1]
		[Total: 13	3]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination project.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The Periodic Table of Elements

	\	2 He	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	첫	krypton 84	54	Xe	xenon 131	98	牊	radon			
	=			6	ш	fluorine 19	17	Cl	chlorine 35.5	35	Ŗ	bromine 80	53	Н	iodine 127	85	¥	astatine -			
	5			80	0	oxygen 16	16	S	sulfur 32	34	Se	selenium 79	52	<u>е</u>	tellurium 128	84	Ъ	polonium –	116		livermorium —
	>			7	z	nitrogen 14	15	₾	phosphorus 31	33	As	arsenic 75	51	Sp	antimony 122	83	Ξ	bismuth 209			
	≥			9	ပ	carbon 12	14	SS	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Ър	lead 207	114	Fl	flerovium —
	=			2	В	boron 11	13	Ρl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	11	thallium 204			
										30	Zu	zinc 65	48	g	cadmium 112	80	£	mercury 201	112	ű	copernicium -
										29	D C	copper 64	47	Ag	silver 108	79	Αn	gold 197	111	Rg	roentgenium -
Group										28	Z	nickel 59	46	Pd	palladium 106	78	చ	platinum 195	110	Ds	darmstadtium -
ğ										27	ပိ	cobalt 59	45	格	rhodium 103	77	'n	indium 192	109	¥	meitnerium —
		- エ	hydrogen 1							26	Ьe	iron 56	44	Ru	ruthenium 101	92	Os	osmium 190	108	Hs	hassium –
										25	Mn	manganese 55	43	ပ	technetium -	75	Re	rhenium 186	107	Bh	bohrium —
				_	pol	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
			Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	q	niobium 93	73	<u>Б</u>	tantalum 181	105	ОР	dubnium —
					atc	re				22	j	titanium 48	40	Zr	zirconium 91	72	士	hafnium 178	104	껖	rutherfordium —
										21	Sc	scandium 45	39	>	yttrium 89	57–71	lanthanoids		89–103	actinoids	
	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	26	Ba	barium 137	88	Ra	radium —
	_			က	=	lithium 7	11	Na	sodium 23	19	×	potassium 39	37	&	rubidium 85	55	CS	caesium 133	87	Ψ̈́	francium —

7.1	P	lutetium	175	103	۲	lawrencium	ı
	Υp	^				_	
69	Tm	thulium	169	101	Md	mendelevium	ı
89	ш	erbinm	167	100	Fm	ferminm	1
29	웃	holmium	165	66	Es	einsteinium	ı
99	۵	dysprosium	163	86	ర్	californium	ı
65	q	terbium	159	97	BK	berkelium	1
64	Gd	gadolinium	157	96	Cm	curium	ı
63	Еn	europium	152	92	Am	americium	ı
62	Sm	samarium	150	94	Pn	plutonium	ı
61	Pm	promethium	I	93	d d	neptunium	I
09	pZ	neodymium	144	92	\supset	uranium	238
69	Ā	praseodymium	141	91	Ра	protactinium	231
28	Ce	cerium	140	06	┖	thorium	232
22	Гa	lanthanum	139	68	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.)