



## Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

CHEMISTRY		0620/43
CENTRE NUMBER	CANDIDATE NUMBER	
CANDIDATE NAME		

Paper 4 Theory (Extended)

May/June 2016

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.


At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



1 The diagram shows a blast furnace.



(a) The following equations represent reactions which take place in the blast furnace.

$$\textbf{A} \quad \textbf{C} \, + \, \textbf{O}_2 \, \rightarrow \, \textbf{CO}_2$$

$$\mathbf{B} \quad \mathsf{CaCO}_3 \, \rightarrow \, \mathsf{CaO} \, + \, \mathsf{CO}_2$$

**D** 
$$CO_2 + C \rightarrow 2CO$$

**E** 
$$Fe_2O_3 + 3CO \rightarrow 2Fe + 3CO_2$$

- (i) Which reaction is used to increase the temperature inside the blast furnace? ............ [1]
- (ii) Which reaction is an example of thermal decomposition? ........... [1]
- (iii) In which reaction is carbon both oxidised and reduced? ............ [1]
- (iv) Which equation shows the removal of an impurity from the iron? ........... [1]
- (v) Which equation shows the reaction of an acidic substance with a basic substance?

......... [1]

**(b)** Use the diagram of the blast furnace to help you answer these questions.

(i) What enters the blast furnace at X?

.....[1]


(ii) What leaves the blast furnace at Y?

\_\_\_\_\_\_[

(iii) Name <b>two</b> waste gases that leave the blast furnace.	(111)	Name <b>two</b> was	ste gases that	leave the b	plast furnace.	
---	-------	---------------------	----------------	-------------	----------------	--

1.	
2.	
	[2]
	[4]

(c) The graph shows how the malleability of iron changes as the percentage of carbon in the iron changes.



(i) Describe how the malleability of iron changes as the percentage of	carbon changes
--	----------------

	[1]
 have ablatical force that black forces a contains binds to related and	


(ii) Iron obtained from the blast furnace contains high levels of carbon.


Explain how the amount of carbon in the iron can be decreased.	


[Total: 12]

2 The structures of six organic compounds are shown.

		`		
H   H—C—   H	-C=	H   =C-	H -C- H	







[2]

[3]

(a) Give the name of F.

© UCLES 2016

[4]
 LL.

**(b)** Identify **two** of the compounds that are members of the same homologous series. Give the general formula of this homologous series.

compounds		 	 	
general form	านla	 	 	

**(c)** Which **two** compounds are isomers of each other? Explain why they are isomers.

compounds	 	 
explanation	 	 

(d) Explain why **B** is an unsaturated hydrocarbon.

0620/43/M/J/16

(e)	Describe how <b>D</b> is manufactured from <b>B</b> . Give a chemical equation for the reaction.
	[3]
(f)	Compound <b>A</b> forms an addition polymer.
	Draw <b>two</b> repeat units of the addition polymer formed from <b>A</b> .

[2]

[Total: 13]

Cle	an dry air contains mainly nitrogen and oxygen.
(a)	Name <b>two</b> other gases that are in clean dry air.
	[2]
<b>/</b> b\	Air often centaine nellutante
(a)	Air often contains pollutants.
	Identify <b>three</b> common gaseous pollutants in air and state how each of these pollutants are produced.
	pollutant gas 1
	how it is produced
	pollutant gas 2
	how it is produced
	pollutant gas 3
	how it is produced
	[6]
	[Total: 8]

(a)	Pot	assium lodide is an ionic compound.
	(i)	Describe what happens, in terms of electron loss and gain, when a potassium atom reacts with an iodine atom.
		[2]
	(ii)	Describe the structure of solid potassium iodide. You may draw a diagram.
		[2]
(	(iii)	Explain why potassium iodide has a high melting point.
		[2]

(b)	Pot	assium iodide and lead nitrate are both soluble. Lead iodide is insoluble.
	(i)	Describe how a pure dry sample of lead iodide could be made from solid potassium iodide and solid lead nitrate.
		[4]
	(ii)	Write an ionic equation for the formation of lead iodide, $PbI_2$ , when potassium iodide and lead nitrate react with each other. State symbols are <b>not</b> required.
		[2]
(c)		en chlorine gas is bubbled through an aqueous solution of potassium iodide, a redox ction takes place. $2I^- + Cl_2 \to I_2 + 2Cl^-$
	/i\	
	(i)	State the colour change expected in this reaction.
		start colour
		end colour[2]
	(ii)	Identify the reducing agent in this reaction. Explain your answer.
		[2]
		[Total: 16]

Dilute hydrochloric acid reacts with sodium carbonate solution.

5

		$2HCl(aq) + Na_2CO_3(aq) \rightarrow 2NaCl(aq) + H_2O(I) + CO_2(g)$
(a)	Exp	plain why effervescence is seen during the reaction.
		[1]
(b)	Dilu	te hydrochloric acid was titrated with sodium carbonate solution.
		<ul> <li>10.0 cm³ of 0.100 mol/dm³ hydrochloric acid were placed in a conical flask.</li> <li>A few drops of methyl orange indicator were added to the dilute hydrochloric acid.</li> <li>The mixture was titrated with sodium carbonate solution.</li> <li>16.2 cm³ of sodium carbonate solution were required to react completely with the acid.</li> </ul>
	(i)	What colour would the methyl orange indicator be in the hydrochloric acid?
		[1]
	(ii)	Calculate how many moles of hydrochloric acid were used.
		mol [1]
(	(iii)	Use your answer to <b>(b)(ii)</b> and the equation for the reaction to calculate the number of moles of sodium carbonate that reacted.
		mol [1]
(	(iv)	Use your answer to <b>(b)(iii)</b> to calculate the concentration of the sodium carbonate solution in mol/dm <sup>3</sup> .
		mol/dm³ [2]
(c)	In a	nother experiment, 0.020 mol of sodium carbonate were reacted with excess hydrochloric d.
		culate the maximum volume (at r.t.p.) of carbon dioxide gas that could be made in this ction.
		dm³ [3]
		[Total: 9]

**6** Concentrated ammonia solution gives off ammonia gas. Concentrated hydrochloric acid gives off hydrogen chloride gas. Ammonia, NH<sub>3</sub>, and hydrogen chloride, HC*l*, are both colourless gases. Ammonia reacts with hydrogen chloride to make the white solid ammonium chloride.

Apparatus is set up as shown.

cotton wool					cotton wool
soaked in concentrated					soaked in concentrated
hydrochloric acid					ammonia solution
	A	В	Ç	D	
27					7—
	glass tu	ıbe			

After ten minutes a white solid forms in the tube where the gases meet.

(a) (i)	Write the chemical equation for the reaction of ammonia with hydrogen chloride.
	[1]
(ii)	Name the process by which the ammonia and hydrogen chloride gases move in the tube.
	[1]
(iii)	At which point, <b>A</b> , <b>B</b> , <b>C</b> or <b>D</b> , does the white solid form? Explain why the white solid forms at that point.
	the solid forms at
	explanation
	[3]
(iv)	The experiment was repeated at a higher temperature.
	Predict how the results of the experiment would be different. Explain your answer.

(b)	Some of the	white solid is	removed from	the tube and	l dissolved in	water.
\ N		WILL DOMA 10		tile tabe alla		Water.

Describe how the white solid could be tested to show it contains,


(i	ammo	

test				
result		 		
	 	 		[3

(ii) chloride ions.

result	
	[3]

**(c)** The diagram shows the electron arrangement in a molecule of ammonia, showing only outer shell electrons.




(i) State the type of bonding in ammonia.



(ii) Hydrazine, N<sub>2</sub>H<sub>4</sub>, is another compound of nitrogen and hydrogen.

Complete the diagram to show the electron arrangement in a molecule of hydrazine, showing only outer shell electrons.



[3]

(	d)	N	lylon	and	protei	ins are	both	ı pol	ymers	containii	าg r	nitrogen

,		on and proteins are sear polymers containing mareger.	
	(i)	Name the linkages found in the polymers of nylon and protein.	
			[1]
	(ii)	Describe <b>one</b> difference in the structures of nylon and protein.	
			[1]
(	iii)	What is the general name given to the products of hydrolysis of proteins?	

(e) Suggest the structure of the monomer used to make the polymer shown.

$$\begin{pmatrix} H & O \\ N & \parallel \\ C & \uparrow \\ n \end{pmatrix}$$

[1]

[Total: 22]

## **BLANK PAGE**

## **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The Periodic Table of Elements

		2 He	helium 4	10	Ne	neon 20	18	Ar	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	R	radon			
				6	ட	fluorine 19	17	Cl	chlorine 35.5	35	Ŗ	bromine 80	53	Н	iodine 127	85	Ą	astatine -			
	>			∞	0	oxygen 16	16	S	sulfur 32	34	Se	selenium 79	52	<u>L</u>	tellurium 128	84	Ъ	molonium —	116	_	livermorium _
	>			7	Z	nitrogen 14	15	₾	phosphorus 31	33	As	arsenic 75	51	Sb	antimony 122	83	Ξ	bismuth 209			
	≥			9	ပ	carbon 12	14	S	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Pb	lead 207	114	Εl	flerovium -
	≡			2	Δ	boron 11	13	Αl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	lΤ	thallium 204			
										30	Zn	zinc 65	48	පි	cadmium 112	80	륀	mercury 201	112	S	copernicium -
										59	Cn	copper 64	47	Ag	silver 108	6/	Au	gold 197	111	Rg	roentgenium -
Group	-									28	Ż	nickel 59	46	Pd	palladium 106	78	풉	platinum 195	110	Ds	darmstadtium -
Ö				,						27	ဝိ	cobalt 59	45	格	rhodium 103	77	٦	iridium 192	109	Μ̈́	meitnerium -
		- I	hydrogen 1							26				Ru	ruthenium 101	92	Os	osmium 190	108	Η̈́	hassium
										25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium
				_	pol	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
			Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	14	Q N	niobium 93	73	Б	tantalum 181	105	ОР	dubnium -
					atc	<u>l</u> e				22	j	titanium 48	40	Zr	zirconium 91	72	Έ	hafnium 178	104	짪	rutherfordium -
										21	Sc	scandium 45	39	>	yttrium 89	57–71	lanthanoids		89–103	actinoids	
	=			4	Be	beryllium 9	12	Mg	magnesium 24	20		calcium 40		ഗ്	strontium 88	56	Ва	barium 137	88	Ra	radium
	_			က	=	lithium 7	=	Na	sodium 23	19	¥	potassium 39	37	В	rubidium 85	55	S	caesium 133	87	ᇁ	francium -

71 Lu	lutetium 175	103	۲	lawrencium	ı
70 <b>Yb</b>	ytterbium 173	102	%	nobelium	ı
e9 Tm	thulium 169	101	Md	mendelevium	ı
88 Д	erbium 167	100	Fm	ferminm	I
67 Ho	holmium 165	66	Es	einsteinium	I
e Dy	dysprosium 163	86	ర్	californium	I
es Tb	terbium 159	26	器	berkelium	I
<sup>2</sup> D	gadolinium 157	96	CB	curium	I
e3 Eu	europium 152	92	Am	americium	ı
62 Sm	samarium 150	94	Pu	plutonium	ı
61 Pm	promethium -	93	d N	neptunium	ı
9 PX	neodymium 144	92	$\supset$	uranium	
59 Pr	praseodymium 141	91	Ра	protactinium	231
Se O	cerium 140	06	드	thorium	232
57 <b>La</b>	lanthanum 139	88	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.)