



Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
CHEMISTRY			0620/33
Paper 3 Theory (Cor	·e)		May/June 2017

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.


The number of marks is given in brackets [] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



1 hour 15 minutes

1 (a) The electronic structures of five atoms, A, B, C, D and E, are shown.



Answer the following questions about these atoms. Each atom may be used once, more than once or not at all.

Which atom, A, B, C, D or E,

(i)	is in Group VIII of the Periodic Table,	 [1]
(ii)	is a chlorine atom,	 [1]
iii)	has 17 protons in its nucleus,	 [1]
iv)	is an atom of an element in the same period as carbon,	 [1]
(v)	is an atom of a metal?	 [1]

(b) Complete the table to show the number of electrons, neutrons and protons in the magnesium atom and calcium ion shown.

		number of electrons	number of neutrons	number of protons
²⁶ N	⁄lg	12		
44 20	a ²⁺		24	

[3]

[Total: 8]

2 (a) The table shows the ions present in a 1000 cm³ sample of mineral water.

ion present	formula of ion	mass present in mg/1000 cm³
calcium	Ca ²⁺	52
chloride	C1-	10
hydrogencarbonate	HCO ₃ -	50
magnesium	Mg ²⁺	
sodium	Na⁺	12
sulfate	SO ₄ ²⁻	10
	NO ₃ -	8
	total	150


Answer these questions using the information from the table.

(i)	Calculate the ma	ass of magnesium	ions in the 1000 cm ³	sample of mineral	water.
\-'/					

	mass of magnesium ions = mg	[1]
(ii)	Which negative ion is present in the highest concentration?	
		[1]
(iii)	State the name of the ion NO ₃ ⁻ .	
		[1]
(iv)	Calculate the mass of hydrogencarbonate ions present in 250 cm ³ of this sample.	

(b)		en nitrate ions are warmed with aqueous sodium hydroxide and aluminium foil, ammo	nia
	Des	scribe a test for ammonia gas.	
	test		
	resu	ult	[2]
(c)	The	formulae of some bromides are given.	[-]
		aluminium bromide, AlBr ₃	
		magnesium bromide, MgBr ₂	
		sodium bromide, NaBr	
	Ded	duce the formula for calcium bromide.	
			[1]
(d)	Mol	ten calcium bromide can be electrolysed using inert electrodes.	
	(i)	Predict the products of this electrolysis at	
		the negative electrode (cathode),	
		the positive electrode (anode).	[2]
	(ii)	Graphite electrodes are inert.	
		Give the name of one other substance that can be used to make an inert electrode.	
			[1]
		[Total:	10]


3 The diagram shows part of the structures of three substances, **P**, **Q** and **R**, at room temperature and pressure.



- (a) Describe substances P, Q and R in terms of
 - their bonding,
 - the arrangement of their particles,
 - the motion of their particles.

	[6]

(b) A closed gas syringe contains substance **R**. The syringe is heated in a water bath.



Describe what happens to the volume of substance R in the syringe. The pressure remain constant. Explain your answer in terms of particles.				
Substar	nce P undergoes physical and chemical changes.			
Which t	wo of the following are physical changes? Explain your answer.			
Α	Substance P reacts with concentrated sulfuric acid.			
В	lodine forms when chlorine is added to an aqueous solution of substance	Ρ.		
С	Substance P boils at 1330 °C.			
D	Substance P dissolves easily in water.			
		[3]		
•	· · ·			
State o	ne other use of graphite and explain how this use is related to its structure.			
		[2]		
		[Total: 12]		
	Substan Which t A B C D Graphit Graphit	Substance P undergoes physical and chemical changes. Which two of the following are physical changes? Explain your answer. A Substance P reacts with concentrated sulfuric acid. B Iodine forms when chlorine is added to an aqueous solution of substance IC Substance P boils at 1330 °C. D Substance P dissolves easily in water. Graphite has a giant covalent structure containing layers of carbon atoms. Graphite is used to make inert electrodes for electrolysis. State one other use of graphite and explain how this use is related to its structure.		

4

Iron is 6	extracted from its ore by heating the ore with carbon in a blast furnace.	
(a) (i)	State the name of an ore of iron.	
		[1]
(ii)	In the blast furnace, iron(III) oxide is reduced by carbon monoxide.	
	Explain how the carbon monoxide is formed in the blast furnace.	
		[2]
(iii)	Balance the chemical equation for this reaction.	
	$Fe_2O_3 + 3CO \rightarrowFe +CO_2$	[2]
(iv)	How does this equation show that iron(III) oxide is reduced?	
		[1]
(v)	Calculate the relative formula mass of iron(III) oxide, ${\rm Fe_2O_3}$. Show all your working. Use your Periodic Table to help you.	
	relative formula mass =	[2]

(b)		Iron reacts with hydrochloric acid to form iron(Π) chloride and a gas which 'pops' with a lighted splint.		
	(i)	Identify this gas.		
		[1]		
	(ii)	Suggest a practical method for investigating the rate of this reaction involving collection of the gas.		
		You may include a labelled diagram in your answer.		
		[3]		
(c)	Des	scribe a test for iron(II) ions.		
	test			
	res	ult		
		[2]		
(d)	Giv	e two advantages of recycling steel.		
	1			
	2			
		[2]		
		[Total: 16]		

0620/33/M/J/17

5 Glycolic acid is found in the stalks of sugar-cane plants.

The structure of glycolic acid is shown.

(a)	On the structure shown draw a circle around the carboxylic acid functional group.	[1]
(b)	Give the molecular formula of glycolic acid showing the number of carbon, hydrogen oxygen atoms.	and
		[1]
(c)	Suggest how you could obtain a solution containing glycolic acid from sugar-cane plants.	
, D		
(d)	Nitric acid can oxidise glycolic acid.	
	What is the meaning of the term oxidation?	.

(e) The table shows the properties of some carboxylic acids.

carboxylic acid	number of carbon atoms in one molecule	melting point in °C	boiling point in °C	density in g/cm³
methanoic acid	1	8	101	1.220
ethanoic acid	2	17	118	1.049
propanoic acid	3	-21		0.993
butanoic acid	4	- 5	164	0.958

(i)	Describe how the density of the carboxylic acids varies with the number of carbon atoms in one molecule.
	[1]
(ii)	Predict the boiling point of propanoic acid.
	[1]
(iii)	What is the state of butanoic acid at -10 °C? Explain your answer.
	[2]
	[Total: 10]

6 (a) The table shows the properties of some alloys.

alloy	density in g/cm³	relative hardness	relative strength	relative electrical conductivity	cost
J	7.8	4.0	24.0	1.1	cheap
K	2.8	2.5	7.5	3.8	expensive
L	11.3	0.2	1.5	0.5	cheap
M	10.2	5.5	16.5	0.2	very expensive


Use the information in the table to answer the questions.

(i)	Which alloy would be most useful for making a bridge? Give two reasons for your answer.	
	alloy	
	reason 1	
	reason 2	[2]
(ii)	Which alloy is best to make the tips of high-speed drills? Give one reason for your answer.	
	alloy	
	reason	[1]
(iii)	Which alloy is best to make aircraft bodies? Give one reason for your answer.	
	alloy	
	reason	
		[1]

(b) A student took pieces of four different steel alloys, W, X, Y and Z, each of the same mass, and placed them separately into hydrochloric acid. The concentration of acid was the same in each case and the metal was in excess. All other conditions were kept the same.

The student measured the mass of each alloy at intervals as the reaction proceeded and calculated the percentage mass loss.

The results for alloys **W**, **X** and **Y** are shown on the graph.



(i) Alloy **Z** reacts faster with hydrochloric acid than alloy **W**.

On the graph, draw a line which could represent the percentage mass loss of alloy **Z** with time.

[2]

(ii) Which alloy showed the least percentage mass loss after 3 days?

.....[1]

(iii) How long did it take for alloy **X** to lose 40% of its mass?

.....[1

(iv) Suggest how the following factors affect the rate of mass loss.

increasing the temperature

increasing the concentration of the acid

[2]

(c) The concentration of an acid can be found by titrating it with aqueous sodium hydroxide.


Suggest which **one** of these pH values is the pH of concentrated aqueous sodium hydroxide. Draw a circle around the correct answer.

pH 1 pH 3 pH 7 pH 12 [1]

[Total: 11]

7 Water is a simple covalent compound.





[1]

(b)	Give two physical properties which distinguish a simple covalent compound from an i compound.	onio
	1	
	2	
		[2
(c)	Some information about the reaction of four metals with water is given.	

cerium: reacts slowly with cold water

iron: reacts with steam only when extremely hot

lithium: reacts rapidly with cold water magnesium: reacts slowly with hot water

List these metals in order of their reactivity. Put the least reactive metal first.

least reactive —		→ most reactive

[2]

(d)	(i)	State the conditions needed for iron to rust.	
	(ii)	State two methods of rust prevention.	
		1	
		2	 [2]
(e)		rting with an aqueous solution of copper(II) sulfate, describe how you could obtain a pusample of copper(II) sulfate crystals.	re
		[2]
(f)	Car	bon dioxide and water are formed when hydrocarbons burn.	
	Cor	mplete the chemical equation for the combustion of butene.	
		$C_4H_8 + 6O_2 \rightarrow \dots CO_2 + \dots H_2O$	[2]

[Total: 13]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the line promised to a series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The Periodic Table of Elements

								Gro	Group								
=													≥	>		₹	₹
							- I										2 He
Key	Key	Key	Key				hydrogen 1										helium 4
4 atomic number	atomic number	atomic number	tomic number									2	9	7	8	6	10
Be atomic symbol	atomic symbol	atomic symbol	mic symbol	loc								В	ပ	z	0	ட	Ne
beryllium name 9 relative atomic mass	name relative atomic mass	name relative atomic mass	name live atomic mass	SS								boron 11	carbon 12	nitrogen 14	oxygen 16	fluorine 19	neon 20
12												13	14	15	16	17	18
Mg												Αl	Si	₾	ഗ	Cl	Ā
magnesium 24												aluminium 27	silicon 28	phosphorus 31	sulfur 32	chlorine 35.5	argon 40
21 22 23 24	22 23 24	23 24	24			55	26	27	28	29	30	31	32	33	34	35	36
Sc Ti Cr	, C	ပ် >	ပ်		2	Mn	Fe	ပိ	Z	Cn	Zu	Ga	Ge	As	Se	Ā	궃
calcium scandium titanium vanadium chromium mang 40 45 48 51 52 5	titanium vanadium chromium r 48 51 52	vanadium chromium r 51 52	chromium r		mang	nanganese 55	iron 56	cobalt 59	nickel 59	copper 64	zinc 65	gallium 70	germanium 73	arsenic 75	selenium 79	bromine 80	krypton 84
39 40 41 42	40 41 42	41 42	42			43	44	45	46	47	48	49	50	51	52	53	54
Y Zr Nb Mo	Zr Nb Mo	Nb Mo	Mo			ان.	Ru	뫈	Pd	Ag	ည	In	Sn	Sb	Те	Н	Xe
zirconium niobium molybdenum 91 93 96	zirconium niobium molybdenum 91 93 96	niobium molybdenum 93 96	molybdenum 96		techr	netium -	ruthenium 101	rhodium 103	palladium 106	silver 108	cadmium 112	indium 115	tin 119	antimony 122	tellurium 128	iodine 127	xenon 131
57–71 72 73 74	72 73 74	73 74	74			75	92	77	78	62	80	81	82	83	84	85	98
lanthanoids Hf Ta W	Hf Ta W	Ta W	>		ш	Şe	Os	'n	풉	Αn	БĤ	11	Pb	<u>B</u>	Ъо	¥	R
tantalum tungsten 181 184	tantalum tungsten 181 184	tantalum tungsten 181 184	tungsten 184		£.	rhenium 186	osmium 190	iridium 192	platinum 195	gold 197	mercury 201	thallium 204	lead 207	bismuth 209	molonium –	astatine	radon
89–103 104 105 106	104 105 106	105 106	106			107	108	109	110	111	112		114		116		
actinoids Rf Db Sg	Rf Db Sg	Db Sg	Sg			Bh	Hs	Ĭ	Ds		ű		Εl		_		
	dubnium seaborgium	dubnium seaborgium	seaborgium			oohrium	hassium	meitnerium	damstadtium	9	copernicium		flerovium		livermorium		
		1	_	\dashv					-		ı		_		ı		

71 Lu	lutetium 175	103	۲	lawrencium	I
70 Yb	ytterbium 173	102	%	nobelium	ı
°9 Tm	thulium 169	101	Md	mendelevium	I
₈₈ Ё	erbium 167	100	Fm	fermium	ı
67 Ho	holmium 165	66	Es	einsteinium	ı
% Dy	dysprosium 163	86	ర్	californium	ı
e5 Tb	terbium 159	26	Ř	berkelium	I
Gd 64	gadolinium 157	96	Cm	curium	I
e3 Eu	europium 152	92	Am	americium	ı
ss Sm	samarium 150	94	Pn	plutonium	ı
Pm	promethium -	93	ď	neptunium	ı
9N	neodymium 144	92	\supset	uranium	238
59 Pr	praseodymium 141	91	Ра	protactinium	231
Se Ce	cerium 140	06	드	thorium	232
57 La	lanthanum 139	68	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).