



Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/43

Paper 4 Theory (Extended)

May/June 2018

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.



The following are the symbols and formulae of some elements and compounds.										
		Ar	Ca(OH) ₂	Cl_2	CO ₂	Cu	Fe	SO ₂	V_2O_5	
		_	questions us pound may b					•		
Stat	te which ele	ment o	or compound	is used	d:					
(a)	to kill bacte	eria in d	drinking wate	er						. [1]
(b)	as a food p	reserv	ative	•••••						. [1]
(c)	as an elect	rical co	onductor in c	ables						. [1]
(d)	as an inert	atmos	phere in lam	ps						. [1]
(e)	to neutralis	e exce	ess acidity in	soil						. [1]
(f)	as a catalys	st in th	e Contact pr	ocess.						. [1]

[Total: 6]

2

(a)	²⁹ A		pe of alumin	ium. The only non-ra	adioactive isotope of aluminium is						
	(i)	Describe, in terms of protons, neutrons and electrons, how the isotopes $^{29}\mathrm{A}\mathit{l}$ and $^{27}\mathrm{A}\mathit{l}$ are similar and how they are different.									
		how they are similar									
		how they are differen	t								
					[2]						
	(ii)	Complete the table to ion.	show the nu	umber of nucleons, r	neutrons and electrons in an $^{27}_{13}$ A l^{3+}						
				number in $^{27}_{13}$ A l^{3+}							
			nucleons								
			neutrons								
			electrons								
				<u> </u>	[3]						
(b)	Alu	minium is extracted fro	om its ore by	electrolysis.							
	(i)	Name the main ore o	f aluminium.								
					[1]						
	(ii)	Why is aluminium no	t extracted from	om its ore by reduction	on with carbon?						
					[1]						
((iii)	The main ore of aluminium contains aluminium oxide. Aluminium oxide is dissolved in molten cryolite before it is electrolysed.									
		Give two reasons, ot	her than cost	, why cryolite is used	1 .						
		1									
		2			[2]						

(iv)	The reaction at the anode during the extraction of aluminium by electrolysis is shown.
	$2O^{2-} \rightarrow O_2 + 4e^-$
	Is this process oxidation or reduction? Give a reason for your answer.
(v)	During the extraction of aluminium by electrolysis, carbon dioxide is formed at the anode.
	Explain how carbon dioxide is formed at the anode.
	[2]
	hen a piece of zinc metal is added to $copper(II)$ sulfate solution there is an immediate action.
	$Zn + CuSO_4 \rightarrow ZnSO_4 + Cu$
	nen a piece of aluminium metal is added to $copper(\mathrm{II})$ sulfate solution the initial reaction is y slow.
(i)	Explain why zinc metal reacts with $copper(II)$ sulfate.
	[1]
(ii)	What type of reaction is this?
	[1]
(iii)	Explain why the initial reaction between aluminium metal and copper(II) sulfate is very slow.
	[1]
	[Total: 15]

Cobalt is	a transition element. Potassium is in Group I of the Periodic Table.
(a) State	e one physical property that is similar for cobalt and potassium.
	[1]
(b) (i)	State one physical property that is different for cobalt and potassium.
	[1]
(ii)	Describe how the physical property given in (b)(i) is different for cobalt compared to
	potassium.
	[1]
	en a small piece of potassium is added to cold water, the potassium floats and disappears reacts.
	e two other observations that would be made when a small piece of potassium is added to water.
1	
2	[2]
	alt reacts with dilute hydrochloric acid to make the salt cobalt(Π) chloride. Bubbles of ogen gas are produced.
(i)	Describe a test for hydrogen.
	test
	result
	[2]
	The rate of reaction of cobalt with dilute hydrochloric acid can be made faster by heating the acid or by increasing its concentration.
	State one other way to make the rate of reaction faster.
	[1]
	Use collision theory to explain how heating the dilute hydrochloric acid makes the rate of reaction faster.
	[2]

(e) When $\mathsf{cobalt}(II)$ chloride is added to water an equilibrium is established.

$$[\mathrm{CoC} l_4]^{2-} + 6\mathrm{H_2O} \iff [\mathrm{Co}(\mathrm{H_2O})_6]^{2+} + 4\mathrm{C} l^{-}$$
 blue pink

(i	i)	A student adds water to a blue solution containing $[CoCl_4]^{2-}$ ions.
		Describe what the student observes. Give a reason for your answer in terms of the position of the equilibrium.
		[2]
(ii	i)	Another student cools a blue solution containing $[CoCl_4]^{2-}$. The blue solution turns pink.
		What does this information indicate about the forward reaction?
		[1]
(f) A	\no	ther compound of cobalt is Co(OH) ₃ .
L	ed	uce the charge on the cobalt ion in Co(OH) ₃ .
		[1]
		[Total: 15]


4	Ethanol is	s a member	of the	homologous	series of	alcohols
_		J 4 1110111001	01 1110	1101110104040		aiooiioio

(a)	Give two	characteristics	of	members	of	a homo	logous	series.
۸	\ /			•		•		5	

1	
2	
	[2

(b) The structure of ethanol is shown.

Complete the dot-and-cross diagram to show the electron arrangement in a molecule of ethanol. Show outer shell electrons only.



[2]

(c) Ethanol can be produced by the catalytic addition of steam to ethene or by the fermentation of glucose.

(i)	Write a chemical	equation	for the	production	of	ethanol	by the	catalytic	addition	of	steam
	to ethene.										

(41
a i i i i
r.1

(ii) Write a chemical equation for the production of ethanol by the fermentation of glucose, $C_6H_{12}O_6$.

Г	41

(iii) State **one** advantage of producing ethanol by the catalytic addition of steam to ethene. Your answer must **not** refer to cost.



(iv) State **one** advantage of producing ethanol by the fermentation of glucose. Your answer must **not** refer to cost.

[1]
 Lil

(d)	Eth	anol can be oxidised to ethanoic acid.	
	Sta	te the chemical reagent needed to oxidise ethanol to ethanoic acid.	
		[[1]
(e)		anoic acid reacts with ethanol in the presence of an acid catalyst. The products are a anic compound and water.	an
	(i)	Draw the structure of the organic compound formed. Show all of the atoms and all of the bonds.	те
	(ii)	State the name of the organic compound formed.	[2]
		[[1]
	(iii)	Which homologous series does the organic compound formed belong to?	
(f)	cop	anoic acid, CH ₃ COOH, is a weak acid. It reacts with copper(II) carbonate to form the saper(II) ethanoate, Cu(CH ₃ COO) ₂ .	
	(i)	What is meant by the term <i>weak</i> when applied to acids?	
	(ii)	Describe how a crystalline sample of copper(II) ethanoate can be prepared starting wi ethanoic acid and copper(II) carbonate.	ith
((iii)	Write the word equation for the reaction between ethanoic acid and copper(II) carbonat	
		[[1]


[Total: 18]

5	(a)	Nickel(II) iodide crystals are hydrated. A sample of hydrated nickel(II) iodide crystals has the
		following composition by mass: Ni, 14.01%; I, 60.33%; H, 2.85%; O, 22.81%.

Calculate the empirical formula of the hydrated nickel(II) iodide crystals.

empirical	formula	=	 [2]


(b) Molten nickel(II) iodide can be electrolysed using the apparatus shown.



During electrolysis, charge is transferred through the copper wires and through the molten nickel(II) iodide.

(i)	Name the type of particles which transfer charge through the copper wires.	
		[1]
(ii)	Name the type of particles which transfer charge through the molten nickel(II) iodide.	
		[1]
iii)	Predict the products of the electrolysis of molten nickel(II) iodide. Write an ior half-equation for the formation of $\bf one$ of these products.	nic
	products	
	ionic half-equation	
		[3]

(c) A student electrolysed copper(II) sulfate solution using the two sets of apparatus shown.



In apparatus **A** the student used carbon electrodes. In apparatus **B** the student used copper electrodes.

The student made the following observations.

apparatus A	apparatus B
The mass of the negative electrode increased.	The mass of the negative electrode increased.
The mass of the positive electrode stayed the same.	The mass of the positive electrode decreased.
Bubbles were seen at the positive electrode.	No bubbles were seen at the positive electrode.

(i)	Explain why the mass of the negative electrode increased in both sets of apparatus.	
		[1]
(ii)	Name the gas that formed the bubbles seen in apparatus A .	
		[1]
(iii)	Explain why the mass of the positive electrode decreased in apparatus B .	
		[4]

electrolysis progresses. Explain your answer.
colour of the solution in apparatus A
colour of the solution in apparatus B
explanation
[3]

(iv) Suggest what happens to the colour of the solution in apparatus **A** and apparatus **B** as the

[Total: 13]

6 Calcium chlorate(V), $Ca(ClO_3)_2$, is made by reacting calcium hydroxide with chlorid
--

$$6Ca(OH)_2 + 6Cl_2 \rightarrow Ca(ClO_3)_2 + 5CaCl_2 + 6H_2O$$

		$6Ca(OH)_2 + 6Cl_2 \rightarrow Ca(ClO_3)_2 + 5CaCl_2 + 6H_2O$
(a)	8.8	8g of calcium hydroxide and 7200 cm³ of chlorine gas are mixed together.
	(i)	How many moles is 8.88g of calcium hydroxide?
	(ii)	mol [2] How many moles of chlorine gas is 7200 cm ³ ?
		mol [1]
	(iii)	What is the maximum number of moles of calcium chlorate(V) that can be made from 8.88 g of calcium hydroxide and 7200 cm ³ of chlorine gas?
		mol [1]
	(iv)	What is the maximum mass of calcium chlorate(V) that can be made from 8.88g of calcium hydroxide and 7200 cm³ of chlorine gas?
		e experiment is repeated using different amounts of calcium hydroxide and chlorine gas. It is maximum mass of calcium chlorate (V) that can be made in the experiment is 4.84 g.
	(v)	The actual mass of calcium $\mbox{chlorate}(V)$ made in the experiment is 3.63 g.
		Calculate the percentage yield. percentage yield = % [1]
		percentage yield/// [1]
(b)	Cal	cium chlorate(V) undergoes thermal decomposition.
	The	e only products are calcium chloride and a colourless gas.
	(i)	What must be done to calcium $\operatorname{chlorate}(V)$ to make it thermally decompose?
		[1]
	(ii)	Write a chemical equation for the thermal decomposition of calcium chlorate (V) .

(c)	Chl	$oric(V)$ acid, $HClO_3$, is a strong acid. It can be made from calcium chlorate(V).	
	(i)	What colour is methyl orange indicator in chloric(V) acid?	
			[1]
	(ii)	Define the term acid in terms of proton transfer.	
			[1]
((iii)	Complete the chemical equation to show $\mathrm{HC}l\mathrm{O_3}$ behaving as an acid in water.	
		$HClO_3 + H_2O \rightarrow \dots + \dots + \dots$	[4]
			[1]
			[Total: 13]

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The Periodic Table of Elements

	110.7		ه ح ل	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	牊	radon			
	18.7				6	ш	fluorine 19	17	Cl	chlorine 35.5	35	ğ	bromine 80	53	П	iodine 127	85	Ą	astatine -			
	3	>			80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	Тe	tellurium 128	84	Po	molodium –	116		livermorium -
	;	>			7	z	nitrogen 14	15	۵	phosphorus 31	33	As	arsenic 75	51	Sb	antimony 122	83	<u>B</u>	bismuth 209			
	2	≥			9	O	carbon 12	14	:S	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Pb	lead 207	114	Εl	flerovium -
	=	=			2	М	boron 11	13	Ν	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	<i>1</i> 1	thallium 204			
											30	Zu	zinc 65	48	р С	cadmium 112	80	Hg	mercury 201	112	S	copernicium -
											59	Cn	copper 64	47	Ag	silver 108	62	Au	gold 197	111	Rg	roentgenium -
	eroup										28	Ż	nickel 59	46	Pd	palladium 106	78	₽	platinum 195	110	Ds	darmstadtium -
5	5				7						27	ပိ	cobalt 59	45	格	rhodium 103	77	٦	iridium 192	109	Ĭ	meitnerium -
			- I	hydrogen 1							26	Fe	iron 56	44	Ru	ruthenium 101	92	Os	osmium 190	108	Hs	hassium
								1			25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium
					_	loq	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≥	tungsten 184	106	Sg	seaborgium -
				Kev	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	g	niobium 93	73	<u>a</u>	tantalum 181	105	В	
						atc	re				22	j	titanium 48	40	Zr	zirconium 91	72	茔	hafnium 178	104	¥	rutherfordium -
											21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
	:	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	56	Ba	barium 137	88	Ra	radium
		_			6	=	lithium 7	#	Na	sodium 23	19	×	potassium 39	37	В	rubidium 85	22	Cs	caesium 133	87	ቷ	francium -

71 Lu	lutetium 175	103	ר	lawrencium	ı
70 Yb				_	
mL Tm	thulium 169	101	Md	mendelevium	ı
₈₈ П	erbium 167	100	Fm	ferminm	ı
67 Ho	holmium 165	66	Es	einsteinium	ı
® Dy	dysprosium 163	86	ŭ	californium	ı
e5 Tb	terbium 159	26	Ř	berkelium	ı
² PO	gadolinium 157	96	Cm	curium	ı
e3 Eu	europium 152	92	Am	americium	ı
62 Sm	samarium 150	94	Pu	plutonium	ı
e1 Pm	promethium	93	δ	neptunium	ı
P N	neodymium 144	92	\supset	uranium	238
59 P	praseodymium 141	91	Ра	protactinium	231
Se Oe	cerium 140	06	Ч	thorium	737
57 La	lanthanum 139	88	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is $24\,dm^3$ at room temperature and pressure (r.t.p.).