



Cambridge Assessment International Education

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
CHEMISTRY			0620/62
Paper 6 Alterna	ative to Practical		May/June 2019
			1 hour
Candidates ans	swer on the Question Paper.		

READ THESE INSTRUCTIONS FIRST

No Additional Materials are required.

Write your centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This syllabus is regulated for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate. This document consists of 8 printed pages and 4 blank pages.



A st	tude	nt did the following steps to make zinc chloride crystals from solid zinc oxide.	
ste ste ste	p 2 p 3	Pour 40 cm³ of dilute hydrochloric acid into a beaker. Add a small amount of zind Warm the mixture and stir it. Continue to add zinc oxide to the beaker until all of the dilute hydrochloric a reacted. Remove the excess zinc oxide. Obtain crystals of zinc chloride from the solution.	
(a)	Nar	me the apparatus used in step 1 to:	
	(i)	add the zinc oxide	
			[1]
	(ii)	warm the mixture.	
			[1]
(b)	Hov	w did the student know that all of the dilute hydrochloric acid had reacted in step 2?	
(c)	(i)	What is meant by the term excess in step 3?	[1]
	(ii)	How is the excess zinc oxide removed in step 3 ?	
			[1]
(d)		scribe how the crystals are obtained in step 4 .	
(e)	Sug	ggest how the method would differ if zinc carbonate were used instead of zinc oxide) .
			[1]
		Γ	Total: 9

2 A student investigated the rate of reaction between magnesium ribbon and solutions of dilute hydrochloric acid of different concentrations, solutions **H**, **I**, **J** and **K**. The dilute hydrochloric acid was in excess in all experiments.

Five experiments were done.

Experiment 1

- A measuring cylinder was used to pour 30 cm³ of solution **H** into a beaker.
- A 5.0 cm length of magnesium ribbon was then added to the beaker.
- A timer was started immediately.
- The time taken for all of the magnesium ribbon to react and to disappear completely was measured.

Experiment 2

Experiment 1 was repeated but using solution I instead of solution H.

Experiment 3

Experiment 1 was repeated but using solution J instead of solution H.

Experiment 4

- Experiment 1 was repeated but using solution **K** instead of solution **H**.
- (a) Use the stop-clock diagrams to record the time taken for each experiment in the table.


experiment	solution	concentration of hydrochloric acid in mol/dm³	stop-clock diagram	time taken for the magnesium ribbon to disappear completely/s
1	н	2.0	seconds 45 15 15 15 minutes	
2	I	1.5	45 15 5 15	
3	J	1.0	45 15 5 15	
4	К	0.8	45 15 5 15	

Experiment 5

Solution **J** was added to some magnesium ribbon in a test-tube. The gas produced was tested. The observations were recorded in the table.

	rapid effervescence and the test-tube felt hot
observations	lighted splint 'popped'

(b) Plot the results for Experiments 1–4 on the grid. Draw a smooth line graph.



time taken for the magnesium ribbon to disappear completely/s

(c) From your graph, deduce the time taken for the magnesium ribbon to disappear completely if a solution of hydrochloric acid of concentration 2.5 mol/dm³ were used.

er.
ć

.....[3

(d)	(i)	i) Why was the same length of magnesium used in Experiments 1–4?	
	(ii)	Suggest the effect on the results if Experiments 1–4 were repeated using 2.5 cm lengths of magnesium ribbon instead of 5.0 cm lengths of magnesium ribbon. Explain your answer.	
		[1]	
(e)	bet\ wou	igest a different method which a student could use to investigate the rate of reaction ween magnesium ribbon and dilute hydrochloric acid. State the apparatus the student ild use and the measurements the student would take.	
		aratus	
	mea	asurements	
		[3]	
(f)	Use	the observations from Experiment 5 to answer these questions.	
	(i)	What type of chemical reaction occurs when magnesium ribbon reacts with dilute hydrochloric acid?	
		[1]	
	(ii)	Identify the gas produced[1]	
		[Total: 16]	

J		Tests were done on solid L and solid M .			
	tests on solid L				
	Com	plete the expected observations.			
	(a)	Describe the appearance of solid L .			
		[1]			
	Solid	d L was divided into two portions.			
		The first portion of solid ${\bf L}$ was heated in a hard-glass test-tube. Any gas produced was tested with cobalt(II) chloride paper.			
		observations			
		[3]			
		second portion of solid ${\bf L}$ was added to distilled water. The mixture was shaken to dissolve ${\bf L}$ and form solution ${\bf L}$. The solution of ${\bf L}$ was divided into two equal portions in two test-tubes.			
		An excess of aqueous sodium hydroxide was added to the first portion of solution ${\bf L}$. The mixture was heated and the gas produced was tested.			
		observations			

(d) Dilute nitric acid and aqueous barium nitrate were added to the second portion of solution L.

tests on solid M

Some of the tests and observations are shown.

tests on solid M	observations
Solid M was dissolved in water. The solution was divided into three portions.	
test 1	
An excess of aqueous sodium hydroxide was added to the first portion of the solution.	red-brown precipitate formed
test 2	
An excess of aqueous ammonia was added to the second portion of the solution.	red-brown precipitate formed
test 3	
Dilute nitric acid and aqueous silver nitrate were added to the third portion of the solution.	white precipitate formed

(e)	Identify solid M.	
		[2]

[Total: 9]

4 Azurite is an ore of copper which contains copper(II) carbonate. Azurite contains no other metal ions.

Plan an experiment to show how a sample of copper could be obtained from large lumps of azurite.

Your answer should include:

- descriptions of the reactions involved
- the expected observations.

ou are provided with a large lump of azurite and common laboratory chemicals and apparatus.	
Γ	

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.