

Cambridge IGCSE[™]

| CANDIDATE NAME | | | | | |
|-------------------|--|--|---------------------|--|--|
| CENTRE NUMBER | | | CANDIDATE NUMBER | | |

CHEMISTRY 0620/51

Paper 5 Practical Test May/June 2020

1 hour 15 minutes

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS


- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].
- Notes for use in qualitative analysis are provided in the question paper.

| For Examiner's Use | | |
|--------------------|--|--|
| 1 | | |
| 2 | | |
| 3 | | |
| Total | | |

This document has **12** pages. Blank pages are indicated.



1 You are going to investigate the reaction between dilute hydrochloric acid and two different aqueous solutions of sodium carbonate labelled solution **E** and solution **F**.

Read all of the instructions carefully before starting the experiments.

Instructions

You are going to do three experiments.

(a) Experiment 1

- Fill the burette up to the 0.0 cm³ mark with dilute hydrochloric acid.
- Use the measuring cylinder to pour 25 cm³ of solution **E** into the conical flask.
- Add five drops of thymolphthalein indicator to the conical flask.
- Slowly add dilute hydrochloric acid from the burette to the conical flask, while swirling the flask, until the solution just changes colour.
- Record the burette readings in the table and complete the table.

| | Experiment 1 |
|--|--------------|
| final burette reading/cm ³ | |
| initial burette reading/cm ³ | |
| volume of dilute hydrochloric acid added/cm³ | |

Experiment 2

- Empty the conical flask and rinse it with distilled water.
- Refill the burette with dilute hydrochloric acid.
- Repeat Experiment 1 using five drops of methyl orange indicator instead of thymolphthalein indicator.
- Record the burette readings in the table and complete the table.

| | Experiment 2 |
|--|--------------|
| final burette reading/cm³ | |
| initial burette reading/cm ³ | |
| volume of dilute hydrochloric acid added/cm ³ | |

Experiment 3

- Empty the conical flask and rinse it with distilled water.
- Refill the burette with dilute hydrochloric acid.
- Use the measuring cylinder to pour 25 cm³ of solution **F** into the conical flask.
- Add five drops of methyl orange indicator to the conical flask.
- Slowly add dilute hydrochloric acid from the burette to the conical flask, while swirling the flask, until the solution just changes colour.
- Record the burette readings in the table and complete the table.

| | | | | Experiment 3 | |
|-----|------|------|--|--|-------------------|
| | | | final burette reading/cm³ | | |
| | | | initial burette reading/cm³ | | |
| | | | volume of dilute hydrochloric acid added/cm ³ | | |
| | | | | | [5] |
| (b) | (i) | Wha | at colour change was observed in the conical flash | k in Experiment 1? | |
| | | from | ı to | | |
| | | | | | [1] |
| | (ii) | Wha | at colour change was observed in the conical flash | k in Experiment 2? | |
| | | from | ı to | | |
| | | | | | [1] |
| (c) | | | e the volumes of dilute hydrochloric acid added in any difference. | Experiment 2 and | Experiment 3. |
| | | | | | |
| | | | | | [2] |
| (d) | | | ne the simplest whole number ratio of volumes ents 1 and 2. | of dilute hydrochlo | oric acid used in |
| | | | ratio Experiment 1:Ex | periment 2 = | [1] |
| (e) | | | lume of dilute hydrochloric acid would be required | The state of the s | s repeated using |

volume = [2]

| (f) | The | conical flask was rinsed with distilled water between each experiment. |
|-----|------|---|
| | (i) | Why was the conical flask rinsed? |
| | | |
| | (ii) | Why does it not matter if a little distilled water is left in the flask after it has been rinsed? |
| | | |
| | | [1] |
| (g) | | e two sources of error in the experiments. For each error suggest an improvement that ld reduce the error. |
| | sou | rce of error 1 |
| | imp | rovement 1 |
| | | |
| | sou | rce of error 2 |
| | imp | rovement 2 |
| | | |

[Total: 18]

You are provided with two solids, solid **G** and solid **H**.

Do the following tests on solid **G** and solid **H**, recording all of your observations at each stage.

tests on solid G

| (a) | | ce about half of solid G in a hard-glass test-tube. Heat the solid gently and then strongly cord your observations. | |
|-----|------|--|-----|
| | | | |
| (b) | (i) | Place the remaining half of solid G in a boiling tube. Add about 10 cm ³ of dilute sulfuric acid to the boiling tube. Test any gas produced. | |
| | | Keep the solution formed for use in (c). | |
| | | Record your observations. | |
| | | | |
| | | | |
| | | | [4] |
| | (ii) | Identify the gas produced in (b)(i). | |
| | | | [1] |
| (c) | into | ave the solution from (b) to settle for three minutes. Carefully pour about half of the solute another boiling tube. this portion add aqueous ammonia slowly until in excess. cord your observations. | ion |
| | | | |
| | | | |
| | | | |
| | | | [3] |
| (d) | Wh | at conclusions can you make about solid G ? | |
| | | | [2] |

tests on solid H

| (e) | Carry out a flame test on solid H . Record your observations. |
|-----|---|
| | [1] |
| | solid ${\bf H}$ to about 10 cm³ of distilled water in a boiling tube. Stopper the boiling tube and shake dissolve solid ${\bf H}$ and form solution ${\bf H}$. |
| (f) | Add about 1cm depth of dilute nitric acid and a few drops of aqueous barium nitrate to solution H . Record your observations. |
| | [1] |
| (g) | Identify solid H . |
| | [2] |
| | [Total: 16] |

3 Cobalt, manganese and nickel are metals. They react with dilute hydrochloric acid to form hydrogen gas.

Plan an investigation to find the order of reactivity of these three metals.

You are provided with:

- samples of each metal
- dilute hydrochloric acid
- common laboratory apparatus.

| Your plan must make it clear how your investigation will be a fair test and how you will use your results to place these metals in order of reactivity. |
|---|
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| |
| [A] |

BLANK PAGE

BLANK PAGE

BLANK PAGE

Notes for use in qualitative analysis Tests for anions

| anion | test | test result |
|---|---|--|
| carbonate (CO ₃ ²⁻) | add dilute acid | effervescence, carbon dioxide produced |
| chloride (C l^-) acidify with dilute nitric acid, then add aqueous silver nitrate | | white ppt. |
| bromide (Br ⁻) [in solution] | acidify with dilute nitric acid, then add aqueous silver nitrate | cream ppt. |
| iodide (I ⁻) [in solution] | acidify with dilute nitric acid, then add aqueous silver nitrate | yellow ppt. |
| nitrate (NO ₃ ⁻) add aqueous sodium hydroxide, then aluminium foil; warm carefully | | ammonia produced |
| sulfate (SO ₄ ²⁻) [in solution] | acidify, then add aqueous barium nitrate | white ppt. |
| sulfite (SO ₃ ²⁻) | add dilute hydrochloric acid, warm gently and test for the presence of sulfur dioxide | sulfur dioxide produced will turn acidified aqueous potassium manganate(VII) from purple to colourless |

Tests for aqueous cations

| cation | effect of aqueous sodium hydroxide | effect of aqueous ammonia | |
|---|---|---|--|
| aluminium (Al³+) | white ppt., soluble in excess, giving a colourless solution | white ppt., insoluble in excess | |
| ammonium (NH ₄ ⁺) ammonia produced on warmin | | - | |
| calcium (Ca ²⁺) white ppt., insoluble in excess | | no ppt., or very slight white ppt. | |
| chromium(III) (Cr ³⁺) | green ppt., soluble in excess | grey-green ppt., insoluble in excess | |
| copper(II) (Cu ²⁺) | light blue ppt., insoluble in excess | light blue ppt., soluble in excess, giving a dark blue solution | |
| iron(II) (Fe ²⁺) | green ppt., insoluble in excess | green ppt., insoluble in excess | |
| iron(III) (Fe ³⁺) | red-brown ppt., insoluble in excess | red-brown ppt., insoluble in excess | |
| zinc (Zn ²⁺) | white ppt., soluble in excess, giving a colourless solution | white ppt., soluble in excess, giving a colourless solution | |

Tests for gases

| | · | |
|-----------------------------------|--|--|
| gas | test and test result | |
| ammonia (NH ₃) | turns damp red litmus paper blue | |
| carbon dioxide (CO ₂) | turns limewater milky | |
| chlorine (Cl ₂) | bleaches damp litmus paper | |
| hydrogen (H ₂) | 'pops' with a lighted splint | |
| oxygen (O ₂) | relights a glowing splint | |
| sulfur dioxide (SO ₂) | turns acidified aqueous potassium manganate(VII) from purple to colourless | |

Flame tests for metal ions

| metal ion | flame colour |
|--------------------------------|--------------|
| lithium (Li ⁺) | red |
| sodium (Na ⁺) | yellow |
| potassium (K ⁺) | lilac |
| copper(II) (Cu ²⁺) | blue-green |

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.