

Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/51

Paper 5 Practical Test

May/June 2021

1 hour 15 minutes

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].
- Notes for use in qualitative analysis are provided in the question paper.

For Examiner's Use		
1		
2		
3		
Total		

This document has 12 pages. Any blank pages are indicated.

1 You are going to investigate the temperature decrease when sodium hydrogencarbonate reacts with dilute hydrochloric acid.

Read all of the instructions carefully before starting the experiments.

Instructions

You are going to do six experiments.

(a) Experiment 1

- Use a measuring cylinder to pour 25 cm³ of dilute hydrochloric acid into a conical flask.
- Use a thermometer to measure the initial temperature of the acid. Record the initial temperature in the table.
- Add the 1g sample of sodium hydrogencarbonate to the conical flask. At the same time start the stop-clock.
- Continually stir the acid and sodium hydrogencarbonate mixture in the conical flask using the thermometer.
- Measure the temperature reached by the mixture after 1 minute. Record the temperature
 of the mixture in the table.
- Calculate and record the temperature decrease in the table.
- Rinse the conical flask with distilled water.

Experiment 2

Repeat Experiment 1 using 2g of sodium hydrogencarbonate instead of 1g.

Experiment 3

Repeat Experiment 1 using 3g of sodium hydrogencarbonate instead of 1g.

Experiment 4

Repeat Experiment 1 using 5 g of sodium hydrogenicarbonate instead of 1 g.

Experiment 5

Repeat Experiment 1 using 6g of sodium hydrogencarbonate instead of 1g.


Experiment 6

• Repeat Experiment 1 using 7 g of sodium hydrogencarbonate instead of 1 g.

experiment	mass of sodium hydrogencarbonate/g	initial temperature/°C	temperature after 1 minute/°C	temperature decrease/°C
1	1			
2	2			
3	3			
4	5			
5	6			
6	7			

(b) Complete a suitable scale on the *y*-axis and plot your results from Experiments 1 to 6 on the grid.

Draw **two** best-fit straight lines through your points. The first straight line should be for the first three points and must pass through (0,0). The second straight line should be for the last three points and must be horizontal. Extend your straight lines so that they meet each other.



(c) (i) From your graph, determine the temperature decrease and mass of sodium hydrogencarbonate where your two straight lines meet. Include appropriate units in your answer.

Show clearly on the grid how you worked out your answer.

temperature decrease =	
mass of sodium hydrogencarbonate =	[3]

(ii) Explain why the temperature decrease becomes constant for high masses of sodium hydrogencarbonate.

(d)	The investigation could be repeated with dilute hydrochloric acid of half the concentration, but the same volume.
	Sketch on the grid the graph you would expect to obtain.
	Label your line D . [2]
(e)	Suggest two changes that could be made to the apparatus that would improve the accuracy of the results. For each change explain why it would improve the accuracy of the results.
	change 1
	explanation 1
	change 2
	explanation 2
	[4]

[Total: 19]

You are provided with one solid, solid **E**, and one solution, solution **F**. Do the following tests on the substances, recording all of your observations at each stage.


tests on solid E

(a)	Place about half of solid E in a hard-glass test-tube. Heat the solid gently for about 30 secon Record your observations.	ds.
		[2]
tube		ing
	ce a stopper in the boiling tube and shake the tube to dissolve solid E and form solution E . Ide solution E into three approximately equal portions in two test-tubes and one boiling tube) <u>.</u>
(b)	To the first portion of solution E in a test-tube, add about 1 cm depth of dilute nitric acid follow by a few drops of aqueous barium nitrate. Record your observations.	ved
		[1]
(c)	To the second portion of solution E in a test-tube, add excess aqueous ammonia. Record your observations.	
		[1]
(d)	To the third portion of solution E in the boiling tube, add aqueous sodium hydroxide dropw and then in excess.	ise
	Keep the product for use in (e).	
	Record your observations.	
		[2]
(e)	Gently warm the product from (d) . Test any gas produced. Record your observations.	
		 [1]

(f)	Ider	ntify the three ions contained in solid E .
		[3]
tes	ts or	n solution F
(g)		ry out a flame test on solution F . cord your observations.
		[1]
(h)	Divi	de the remaining solution F into two approximately equal portions in two test-tubes.
	(i)	To the first portion of solution F add a few drops of universal indicator solution. Record your observations.
		[1]
	(ii)	To the second portion of solution ${\bf F}$ add approximately 2cm depth of aqueous copper(II) sulfate. Record your observations.
		[1]
(i)	Ider	ntify solution F .
		[2]

[Total: 15]

3 Dilute hydrochloric acid reacts with calcium carbonate to make carbon dioxide gas. The apparatus shown in the diagram can be used to follow the progress of the reaction. The carbon dioxide gas leaves the flask causing the mass shown on the balance to decrease.



Plan an investigation, using the apparatus shown in the diagram, to find out how the temperature of the dilute hydrochloric acid affects the rate of the reaction. Your plan should include how your results will show how the temperature of the dilute hydrochloric acid affects the rate of the reaction.

You are provided with dilute hydrochloric acid, calcium carbonate and common laboratory apparatu	
F	[6]

BLANK PAGE

BLANK PAGE

BLANK PAGE

Notes for use in qualitative analysis Tests for anions

anion	test	test result
carbonate (CO ₃ ²⁻)	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
bromide (Br ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	cream ppt.
iodide (I ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO ₃ ⁻) [in solution]	add aqueous sodium hydroxide, then aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ²⁻) [in solution]	acidify, then add aqueous barium nitrate	white ppt.
sulfite (SO ₃ ²⁻)	add dilute hydrochloric acid, warm gently and test for the presence of sulfur dioxide	sulfur dioxide produced will turn acidified aqueous potassium manganate(VII) from purple to colourless

Tests for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al³+)	white ppt., soluble in excess, giving a colourless solution	white ppt., insoluble in excess
ammonium (NH ₄ ⁺)	ammonia produced on warming	-
calcium (Ca ²⁺)	white ppt., insoluble in excess	no ppt., or very slight white ppt.
chromium(III) (Cr³+) green ppt., soluble in excess		grey-green ppt., insoluble in excess
copper(II) (Cu²+) light blue ppt., insoluble in excess		light blue ppt., soluble in excess, giving a dark blue solution
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn ²⁺)	white ppt., soluble in excess, giving a colourless solution	white ppt., soluble in excess, giving a colourless solution

Tests for gases

gas	test and test result	
ammonia (NH ₃)	turns damp red litmus paper blue	
carbon dioxide (CO ₂)	turns limewater milky	
chlorine (Cl ₂)	bleaches damp litmus paper	
hydrogen (H ₂)	'pops' with a lighted splint	
oxygen (O ₂)	relights a glowing splint	
sulfur dioxide (SO ₂)	turns acidified aqueous potassium manganate(VII) from purple to colourless	

Flame tests for metal ions

metal ion	flame colour
lithium (Li ⁺)	red
sodium (Na ⁺)	yellow
potassium (K⁺)	lilac
copper(II) (Cu ²⁺)	blue-green

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.