



Cambridge Assessment International Education

Cambridge International General Certificate of Secondary Education

CANDIDATE NAME						
CENTRE NUMBER				CANDIDATE NUMBER		

CHEMISTRY 0620/51

Paper 5 Practical Test

October/November 2019

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Notes for use in qualitative analysis are provided on pages 7 and 8.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
1		
2		
3		
Total		

This syllabus is regulated for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of 8 printed pages.



1 You are going to investigate the reaction between dilute hydrochloric acid and three different concentrations of aqueous sodium hydroxide, labelled **R**, **S** and **T**.

Read all the instructions carefully before starting the experiments.

Instructions

You are going to do three experiments.

Experiment 1

- Fill the burette up to the 0.0 cm³ mark with dilute hydrochloric acid.
- Use the measuring cylinder to pour 20 cm³ of solution **R** into the conical flask.
- Add six drops of methyl orange indicator to the conical flask.
- Add dilute hydrochloric acid from the burette to the conical flask, 1.0 cm³ at a time, while swirling the conical flask, until the solution just changes colour.
- Record the burette readings in the table.
- Empty the conical flask and rinse it with distilled water.

Experiment 2

- Repeat Experiment 1 using solution **S** instead of solution **R**.
- Record the burette readings in the table.

Experiment 3

- Repeat Experiment 1 using solution T instead of solution R.
- Record the burette readings in the table.
- (a) Complete the table.

burette reading/cm ³	Experiment 1 using solution R	Experiment 2 using solution S	Experiment 3 using solution T
final burette reading			
initial burette reading			
volume used			

		[4]
(b)	What colour change is observed in the conical flask at the end-point?	
	from to	[2]
(c)	Suggest why Universal Indicator is not a suitable indicator to use in these experiments.	
		[1]

(d)	(i)	Complete the sentences.
		Experiment needed the smallest volume of dilute hydrochloric acid to change the colour of the methyl orange indicator.
		Experiment needed the largest volume of dilute hydrochloric acid to change the colour of the methyl orange indicator.
		[1]
	(ii)	Determine the simplest whole number ratio of volumes of dilute hydrochloric acid used in Experiments 1 and 2.
		Experiment 1 Experiment 2 [1]
((iii)	Deduce the order of concentrations of the solutions of aqueous sodium hydroxide, ${\bf R},{\bf S}$ and ${\bf T}.$
		most concentrated
		least concentrated
		[1]
(e)		at would be the effect on the results, if any, if the solutions of aqueous sodium hydroxide re warmed before adding the dilute hydrochloric acid? Give a reason for your answer.
	effe	ect on the results
	rea	son[2]
		[2]
(f)	Sug	ggest how the reliability of the results could be checked.
		[0]
		[2]
(g)		ggest a different method, not involving an indicator, of finding the order of concentrations of solutions of aqueous sodium hydroxide, ${f R}$, ${f S}$ and ${f T}$.
		[3]
		[Total: 17]

You are provided with solid U and liquid V.
Do the following tests on solid U and liquid V, recording all of your observations at each stage.

tests on solid U

(a)	Do a flame test on a small sample of solid U . Record your observations.							
			[1]					
		rest of solid ${\bf U}$ to about 10 cm³ of distilled water in a boiling tube. Stopper the boiling tuke the mixture to dissolve solid ${\bf U}$ and form solution ${\bf U}$.	be					
(b)	Des	cribe the colour of solution U .						
			[1]					
Div	ide s	olution U into three approximately equal portions in three test-tubes.						
(c)	(i)	Add a few drops of aqueous sodium hydroxide to the first portion of solution ${\bf U}$. Record your observations.						
			[1]					
	(ii)	Now add an excess of aqueous sodium hydroxide to this mixture. Record your observations.						
			[1]					
(d)	(i)	Add a few drops of aqueous ammonia to the second portion of solution U . Record your observations.						
			[2]					
	(ii)	Now add an excess of aqueous ammonia to this mixture. Record your observations.						
			[1]					
(e)	port	a few drops of dilute nitric acid and about $1\mathrm{cm^3}$ of aqueous barium nitrate to the th ion of solution \mathbf{U} . Leave to stand for 5 minutes. ord your observations.	ird					
			[2]					
(f)	Soli	d U contains three different ions.						
	Wha	at conclusions can you draw about the ions present in solid U ?						

tests on liquid V

(g)	Describe the appearance of liquid V . Record your observations.
	[1]
(h)	Use a teat pipette to place a few drops of liquid V onto a watch-glass. Put the stopper back into the test-tube of liquid V . Use a lighted splint to touch the surface of liquid V carefully. Record your observations.
	[1]
(i)	Use a spatula to transfer a small crystal of iodine carefully into the rest of liquid ${\bf V}$ in the stoppered test-tube. Put the stopper back into the test-tube and shake the test-tube. Record your observations.
	[2]
(j)	Draw one conclusion about liquid V .
	[1]
	[Total: 17]

3 Potassium nitrate and ammonium chloride are two salts. The energy change when they each dissolve in water is endothermic.

Plan an experiment to show which of these two salts produces the larger endothermic energy change per gram.

Your answer should include:

- any measurements you would take and record how the results could be used to draw a conclusion.

You are prov laboratory app	paratus.					
		•••••	 	 •••••	 	[6]

Notes for use in qualitative analysis Tests for anions

anion	test	test result
carbonate (CO ₃ ²⁻)	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
bromide (Br ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	cream ppt.
iodide (I ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO ₃ ⁻) [in solution]	add aqueous sodium hydroxide, then aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ²⁻) [in solution]	acidify, then add aqueous barium nitrate	white ppt.
sulfite (SO ₃ ²⁻)	add dilute hydrochloric acid, warm gently and test for the presence of sulfur dioxide	sulfur dioxide produced will turn acidified aqueous potassium manganate(VII) from purple to colourless

Tests for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia		
aluminium (Al³+)	white ppt., soluble in excess, giving a colourless solution	white ppt., insoluble in excess		
ammonium (NH ₄ ⁺)	ammonia produced on warming	-		
calcium (Ca ²⁺)	white ppt., insoluble in excess	no ppt., or very slight white ppt.		
chromium(III) (Cr³+) green ppt., soluble in excess		grey-green ppt., insoluble in excess		
copper(II) (Cu ²⁺) light blue ppt., insoluble in excess		light blue ppt., soluble in excess, giving a dark blue solution		
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess		
iron(III) (Fe³+) red-brown ppt., insoluble in excess		red-brown ppt., insoluble in excess		
zinc (Zn ²⁺)	white ppt., soluble in excess, giving a colourless solution	white ppt., soluble in excess, giving a colourless solution		

Tests for gases

gas	test and test result			
ammonia (NH ₃)	turns damp red litmus paper blue			
carbon dioxide (CO ₂)	turns limewater milky			
chlorine (Cl ₂)	bleaches damp litmus paper			
hydrogen (H ₂)	'pops' with a lighted splint			
oxygen (O ₂)	relights a glowing splint			
sulfur dioxide (SO ₂)	turns acidified aqueous potassium manganate(VII) from purple to colourless			

Flame tests for metal ions

metal ion	flame colour
lithium (Li ⁺)	red
sodium (Na ⁺)	yellow
potassium (K ⁺)	lilac
copper(II) (Cu ²⁺)	blue-green

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.