

Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/41

Paper 4 Theory (Extended)

October/November 2021

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

1 Some elements are shown in the order they appear in the reactivity series. The most reactive element is at the top.

sodium
calcium
magnesium
aluminium
zinc
iron
hydrogen
copper

		copper	
(a)		swer the questions using the list of elements. Each element may be used once, more that be or not at all.	an
	Ide	ntify:	
	(i)	a non-metal	
	(ii)	a metal which is stored under oil	[1]
		[[1]
((iii)	the main component of steel	
		[[1]
((iv)	a metal with three electrons in the outer shell of its atoms	
		[[1]
	(v)	a metal found in brass	
		[[1]
	(vi)	a metal that forms chlorides of the type $\mathrm{XC}\mathit{l}_{2}$ and $\mathrm{XC}\mathit{l}_{3}$.	
		[[1]
(b)	Nar	me the main ores of:	
	(i)	zinc[[1]
	(ii)	aluminium. [[1]
(c)	In a	an experiment, a sample of aluminium appeared less reactive than expected.	
	Exp	plain why.	
		[[1]

(d)		ne two metals from the list which are extracted by reduction of their ores using carbon.
	2	[2]
(e)		en zinc granules are added to aqueous copper(II) sulfate, a reaction occurs. During the ction, a red-pink solid is formed and the solution becomes colourless.
	(i)	Name the red-pink solid.
	(ii)	Name the colourless solution. [1]
		[1]
(iii)	Explain, in terms of particles, why the rate of this reaction increases when the temperature is increased.
		[3]
(iv)	Suggest two other ways of increasing the rate of this reaction.
		1
		2[2]
		[Total: 18]

0620/41/O/N/21

© UCLES 2021

This question is about copper and its compounds.

2

(a) Co	pper has two diffe	erent naturally	occurring	atoms, ⁶	⁶³ Cu and ⁶⁵ C	Gu.	
(i)	State the term u	sed for atoms	of the sa	me elem	ent with diffe	erent nucleon r	numbers.
							[1]
(ii)	The atomic num	ber of copper	is 29.				
	Complete the ta of copper shown		e number	of proto	ns, neutrons	and electrons	in the particles
			63 C	u	⁶⁵ Cu ²⁺		
		protons					
		neutrons					
		electrons					
						J	[3]
(iii)	Relative atomic	mass is the av	/erage ma	ass of na	aturally occu	rring atoms of	an element.
	The percentage	of the natural	y occurrii	ng atoms	s in a sample	e of copper is s	hown.
			⁶³ Cu	⁶⁵ Cu			
			70%	30%			
	Deduce the rela	tive atomic ma	ass of cop	per in th	is sample.		
	Give your answ	er to one decir	mal place				
				relati	ive atomic m	nass =	[2]

(b)		hydrous copper(II) sulfate is used to test for the presence of water. When this test is posit drated copper(II) sulfate is formed.	ive,
	(i)	State the colour change seen during this test.	
		from to	[2]
	(ii)	Complete the chemical equation to show the reaction that takes place.	
		$CuSO_4 + \dots \rightleftharpoons CuSO_4 • 5H_2O$	[1]
((iii)	State how hydrated copper(II) sulfate can be turned back into anhydrous copper(III) sulfate can be turned back into anhydrous copper(IIII) sulfate can be turned back into anhydrous copper(IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII	ate.
			[1]
((iv)	Describe a test for pure water.	
			[2]
(c)	Aqı	ueous copper(II) sulfate contains Cu²+(aq) ions.	
	(i)	Describe what is seen when aqueous copper(II) sulfate is added to aque sodium hydroxide, NaOH(aq).	ous
			[1]
	(ii)	Write the ionic equation for the reaction between aqueous copper(II) sulfate and aqueout sodium hydroxide.	ous
		Include state symbols.	
			[3]

	6
(d)	When solid copper(II) nitrate is heated copper(II) oxide, nitrogen dioxide and oxygen are formed.
	$2Cu(NO_3)_2 \rightarrow 2CuO + 4NO_2 + O_2$
	Calculate the volume of nitrogen dioxide formed at room temperature and pressure when $4.7\mathrm{g}$ of $\mathrm{Cu(NO_3)_2}$ is heated.
	Use the following steps:
	• calculate the mass of one mole of Cu(NO ₃) ₂
	 calculate the number of moles of Cu(NO₃)₂ used moles determine the number of moles of nitrogen dioxide formed
	moles

calculate the volume of nitrogen dioxide formed at room temperature and pressure.

(e) Write the chemical equation to show the action of heat on sodium nitrate, NaNO₃.


..... dm³

[4]

[Total: 22]

3 This question is about electrolysis.

Concentrated hydrochloric acid is electrolysed using the apparatus shown.



- (a) Chloride ions are discharged at the anode.
 - (i) Complete the ionic half-equation for this reaction.

......
$$Cl^{-}(aq) \rightarrow(g) +e^{-}$$
 [2]

(ii) State whether oxidation or reduction takes place. Explain your answer.

 	 [1]

(b) Describe what is seen at the cathode.

[1]

(c) Write the ionic half-equation for the reaction at the cathode.

FO1
コンロ
 141

(d) The pH of the electrolyte is measured throughout the experiment.

(i)	Suggest the p	oH of the el	ectrolyte at tl	he beginning	of the experime	nt.
-----	---------------	--------------	-----------------	--------------	-----------------	-----

Г1	1
l'	1

(ii) State how the pH changes, if at all, during the experiment.

Explain your answer.

.....[2]

(e)	The electrolysis is repeated using molten lead(II) bromide.
	Describe what is seen at the:
	cathode
	• anode[2]
(f)	State two properties of graphite (carbon) which make it suitable for use as an electrode.
	1
	2
	[2]
	[Total: 13]

- 4 Chalcopyrite, FeCuS₂, is used in the manufacture of sulfuric acid in the Contact process.
 - (a) In the first stage of the process, chalcopyrite reacts with oxygen in the air to produce sulfur dioxide, SO₂, iron(III) oxide and copper(II) oxide.

Complete the chemical equation for the reaction of FeCuS₂ with oxygen.

$$4FeCuS_2 + 13O_2 \rightarrow \dots + \dots + \dots + \dots$$
 [2]

(b) Sulfur dioxide is then converted to sulfur trioxide.

$$2SO_2 + O_2 \rightleftharpoons 2SO_3$$

The reaction is exothermic. It is also an equilibrium.

(i)	State tw	o features	of an	equilibrium.
-----	----------	------------	-------	--------------

1	
2	
	[2]

- (ii) State the temperature and pressure used in this reaction. Include units.
- (iii) Name the catalyst used.

[1]

(iv) Explain why a catalyst is used.

(v) Describe and explain, in terms of equilibrium, what happens when the temperature is increased.

(2)	

(c) Concentrated sulfuric acid is a dehydrating agent.

When glucose is dehydrated, carbon and one other product are formed.

Complete the equation to show the dehydration of glucose, C₆H₁₂O₆.

$$C_6H_{12}O_6 \rightarrowC +$$
 [2]

[Total: 12]

5

	s and carboxylic acids are both families of similar compounds with similar chemical properties. s and carboxylic acids have different reactions.
(a) Sta	te the term used for a 'family' of similar compounds.
	[1]
(b) Sta	te the general formula of alkenes.
	[1]
(c) The	e structure of but-2-ene is shown.
	H H H
(i)	But-2-ene reacts with aqueous bromine in an addition reaction.
	Describe the colour change seen when but-2-ene is added to aqueous bromine.
	from to
(ii)	State what is meant by the term addition reaction.
	[1]
(iii)	Write the chemical equation for the reaction between but-2-ene and bromine.
	[2]
(iv)	But-2-ene forms a polymer.
	Suggest the name of the polymer formed from but-2-ene.
	[1]
(v)	Name and draw a structural isomer of but-2-ene.
	Show all of the atoms and all of the bonds.
	name

© UCLES 2021 0620/41/O/N/21

structure

(d)	But	anoic acid, CH ₃ CH ₂ CH ₂ COOH, is a carboxylic acid.
	(i)	Deduce the empirical formula of butanoic acid.
		[1]
	(ii)	Complete the chemical equation for the reaction of butanoic acid and sodium carbonate, $\mathrm{Na_2CO_3}$.
		$2CH_3CH_2COOH + Na_2CO_3 \rightarrow \dots + \dots + \dots + \dots + \dots$ [2]
	(iii)	Butanoic acid reacts with methanol to form an organic compound and water.
		Name the organic compound formed.
		[1]
		Draw the structure of the organic compound formed.
		Show all of the atoms and all of the bonds.

[2]

[Total: 15]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

The Periodic Table of Elements

	110.7		۵ ۲	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	牊	radon			
	10,7	=			6	ш	fluorine 19	17	Cl	chlorine 35.5	35	ğ	bromine 80	53	Н	iodine 127	85	¥	astatine -			
	3	>			80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	Тe	tellurium 128	84	Ъ	molodium –	116		livermorium -
	;	>			7	z	nitrogen 14	15	۵	phosphorus 31	33	As	arsenic 75	51	Sb	antimony 122	83	Ξ	bismuth 209			
	2	≥			9	O	carbon 12	14	:S	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Pb	lead 207	114	Εl	flerovium -
	=	=			2	Ш	boron 11	13	Ν	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	11	thallium 204			
											30	Zu	zinc 65	48	р С	cadmium 112	80	Нg	mercury 201	112	S	copernicium -
											59	Cn	copper 64	47	Ag	silver 108	62	Au	gold 197	111	Rg	roentgenium -
	eroup										28	Ż	nickel 59	46	Pd	palladium 106	78	Ŧ	platinum 195	110	Ds	darmstadtium -
5	5				7						27	ပိ	cobalt 59	45	格	rhodium 103	77	i	iridium 192	109	Ĭ	meitnerium -
			- I	hydrogen 1							26	Fe	iron 56	44	Ru	ruthenium 101	92	Os	osmium 190	108	Hs	hassium
								1			25	Mn	manganese 55	43	ည	technetium -	75	Re	rhenium 186	107	Bh	bohrium
						loq	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≯	tungsten 184	106	Sg	seaborgium -
				Kev	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	g	niobium 93	73	<u>n</u>	tantalum 181	105	В	
						atc	<u>re</u>				22	j	titanium 48	40	Zr	zirconium 91	72	茔	hafnium 178	104	꿆	rutherfordium -
											21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
	:	=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	ഗ്	strontium 88	56	Ba	barium 137	88	Ra	radium
		_			e	=	lithium 7	#	Na	sodium 23	19	×	potassium 39	37	В	rubidium 85	22	S	caesium 133	87	ቷ	francium -

71	Intetium	175	103	ב	lawrencium	I
02 \$	Y D ytterbium	173	102	%	nobelium	ı
69 F	thulium	169	101	Md	mendelevium	1
88 [erbinm	167	100	Fm	ferminm	ı
29	D lyolminm	165	66	Es	einsteinium	I
99	dysprosium	163	86	ర్	californium	I
9 1	terbium	159	26	BK	berkelium	I
4 (gadolinium	157	96	Cm	curium	I
63	europium	152	92	Am	americium	ı
62	Samarium	150	94	Pu	plutonium	ı
61	Dromethium	ı	93	Δ	neptunium	ı
09	DC neodymium	144	92	\supset	uranium	238
59	praseodymium	141	91	Ра	protactinium	231
288	Serium G	140	06	T	thorium	232
22	L'a	139	88	Ac	actinium	ı

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).