

# **Cambridge IGCSE**<sup>™</sup>

CANDIDATE NAME				
CENTRE NUMBER		CANDIDATE NUMBER		

CHEMISTRY 0620/31

Paper 3 Theory (Core)

October/November 2023

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

#### **INSTRUCTIONS**

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

#### **INFORMATION**

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].
- The Periodic Table is printed in the question paper.

1 A list of substances is shown.

ammonium nitrate
carbon monoxide
copper(II) chloride
ethane
ethene
litmus
methane
methyl orange
sodium chloride
sodium sulfate
sulfur dioxide
thymolphthalein

Answer the following questions using only the substances from the list. Each substance may be used once, more than once or not at all.

Give the name of the substance that:

(a)	turns from blue to colourless when an acid is added	
		[1]
(b)	is in many fertilisers	
		[1]
(c)	is a salt which has a negative ion with a charge of 2–	
		[1]
(d)	is a waste gas from digestion in animals	
		[1]
(e)	is a hydrocarbon with a total of five atoms in a molecule	
		[1]
(f)	is a compound of a transition element.	
		[1]

[Total: 6]

**2** (a) Fig. 2.1 shows the distillation apparatus that can be used to separate water from aqueous copper(II) sulfate.




Fig. 2.1

Explain how distillation separates water from aqueous copper(II) sulfate.							

**(b)** Fig. 2.2 shows a fractionating column for separating petroleum into different hydrocarbon fractions.




Fig. 2.2

(i) On Fig. 2.2, draw an **X** inside the column to show where the hydrocarbon with the highest viscosity collects. [1]

(ii)	Name the fraction labelled <b>A</b> in Fig. 2.2.	
		[1]
iii)	State the name of the fraction in Fig. 2.2 which has the lowest boiling point.	
		[1]
iv)	State <b>one</b> use of the bitumen fraction.	
		[1]

[Total: 6]

**3** (a) Table 3.1 shows the average concentrations, in ng/1000 cm³, of air pollutants in four different years.

Table 3.1

	concentration of air pollutant in ng/1000 cm <sup>3</sup>						
year	ammonia	hydrocarbons	oxides of nitrogen	particulates	sulfur dioxide		
2019	10.6	12.0	15.3	30.1	20.5		
2020	11.2	13.0	21.6	28.2	20.0		
2021	14.3	15.2	23.5	26.5	25.0		
2022	15.5	9.0	14.0	25.2	18.2		

(1)	name the pollutant that has the lowest concentration in 2019.
	[1]
(ii)	Name the pollutant that shows a continuous decrease in concentration from 2019 to 2022.
	[1]
(iii)	Calculate the average mass, in ng, of sulfur dioxide in a 250 cm <sup>3</sup> sample of polluted air in 2020.
	mass = ng [1]
(b) (i)	State <b>one</b> source of sulfur dioxide in the atmosphere.
	[1]
(ii)	State <b>one</b> adverse effect of sulfur dioxide in the atmosphere.
	[1]

(iii)	Choose the compound	used to remove	sulfur dioxic	le in flue gas d	esulfurisation.
	Tick (✓) <b>one</b> box.				
		aluminium chlor	ide		
		calcium oxide			
		methane			
		sulfuric acid			F41
(i)			IC:1 -		[1]
(iv)	Hydrochloric acid reacts	s with sodium sui	itite.		
	The products are sodio cobalt(II) chloride pink.	um chloride, sul	fur dioxide	and a liquid v	which turns anhydrous
	Complete the symbol ed	quation for this re	eaction.		
	Na <sub>2</sub> SO <sub>3</sub> +	+ $HCl \rightarrow 2N$	aCl + SO <sub>2</sub>	<u> </u> +	[2
(v)	Name the acidified solu	tion used to test	for sulfur di	oxide gas and	state the observations
	acidified solution				
	observations				
					[2]
(c) Am	monia forms an alkaline	solution in water			
(i)	Give the formula of the	ion that is preser	nt in all alka	line solutions	
(-)		·			[4]
<b>(!!)</b>					[1]
(ii)	Choose from the list the	ph value for an	aikaiine so	iution.	
	Draw a circle around yo	ur chosen answ	er.		
	pH1	pH4	pH7	pH13	[1]
					[Total: 12]

4

Bro	omine is a liquid at room temperature.	
(a)	State <b>two</b> general properties of a liquid.	
	1	
	2	
		[2]
(b)	Fig. 4.1 shows the physical states of bromine.	
	solid liquid bromine	
	bromine bromine gas	
	Fig. 4.1	
	Name the changes of physical states <b>A</b> and <b>B</b> .	
	A	
	В	
		[2]
(c)	Describe liquid bromine and bromine gas in terms of the arrangement and motion particles.	of the
	liquid bromine	
	arrangement	
	motion	
	bromine gas	
	arrangement	
	motion	
		[4]

(d) A sealed gas syringe contains  $80\,\mathrm{cm^3}$  of bromine gas.

e affects the volume of bromine gas in the gas syringe when	State how decreasing the pressure the temperature remains constant.
[1]	
[Total: 9]	

- 5 This question is about metals and metal compounds.
  - (a) Table 5.1 shows some properties of some Group I metals.

Table 5.1

metal	melting point in °C	boiling point in °C	observations on reaction with water	solubility of metal hydroxide in g/dm³ at room temperature
sodium	sodium 98 883		bubbles form rapidly but no flame	
potassium	63	760		1130
rubidium		686	explodes	1980
caesium	29	669	explodes	3860

Use the information in Table 5.1 to predict:

(i)	the melting point of rubidium	[1]
(ii)	the solubility of sodium hydroxide at room temperature	[1]
(iii)	the observations when potassium reacts with water	
		[1]
(iv)	the physical state of caesium at 20 °C. Give a reason for your answer.	
	physical state	
	reason	
		[2]

(b)	<b>(b)</b> Iron is extracted in a blast furnace by reduction of iron(III) oxide, Fe <sub>2</sub> O <sub>3</sub> , with carbon monoxide.								
	Carbon monoxide is produced by the reaction of carbon with carbon dioxide.								
	$C + CO_2 \rightarrow 2CO$								
	(i)	Explain how t	this equation shows that	carbon dioxi	de is reduced.				
							[1]		
	(ii)	Name the t simultaneous	ype of chemical reac ly.	tion where	oxidation and	reduction take	place		
							[1]		
<b>(</b> i	iii)	Calcium carb	onate is added to the bla	ast furnace.					
		The calcium	carbonate undergoes the	ermal decom <sub>l</sub>	position.				
		State the mea	aning of the term therma	al decomposit	ion.				
							[2]		
(c)	Stai	nless steel is	an alloy of iron.						
	(i)		son why alloys are more	e useful than	oure metals.				
	(-)						[1]		
	(ii)	Brass is an a					[.]		
	Choose the diagram, <b>A</b> , <b>B</b> , <b>C</b> or <b>D</b> , in Fig. 5.1 that best shows the structure of brass.								
Cu(z Zn Cu(t	Zn Cu/Z	Zn Cu		Fe Cu) Cu) Fe Cu) Cu)	Zn Zn	Fe Zn Fe Z Fe Fe Fe Zn Fe Fe F	(Zn)		
	A	<b>\</b>	В		С	D			
	Fig. 5.1								

diagram ......[1]

(d) Table 5.2 gives some observations about the reactivity of four metals with dilute hydrochloric acid.

Table 5.2

metal	observations
iron	bubbles form slowly
magnesium	bubbles form very quickly
mercury	no bubbles form
tin	bubbles form very slowly

Put the four metals in order of their reactivity. Put the least reactive metal first.

least reactive -	_		$\dashv$	most reactive

[2]

[Total: 13]

- **6** A student investigates the reaction of large pieces of magnesium carbonate with dilute hydrochloric acid at 20 °C. The magnesium carbonate is in excess.
  - (a) Fig. 6.1 shows the volume of carbon dioxide gas released as the reaction proceeds.




Fig. 6.1

(i) Deduce the volume of carbon dioxide gas released after 2 minutes.

volume of carbon dioxide = ...... cm<sup>3</sup> [1]

(ii) The student repeats the experiment using the same volume of hydrochloric acid but with a higher concentration. The magnesium carbonate is still in excess.

All other conditions stay the same.

Draw a line on the grid in Fig. 6.1 to show the volume of carbon dioxide released when hydrochloric acid with a higher concentration is used. [2]

(b)	(i)	The student	t repea	ats the experimer	nt usin	g smaller pieces o	of mag	gnesium carbonat	e.
		All other co	ndition	s stay the same.					
		Describe housed.	w the	rate of reaction di	iffers v	vhen smaller piece	es of n	nagnesium carbor	nate are
									[1]
	(ii)	The student	t repea	ats the experimer	nt at 1	0°C.			
		All other co	ndition	s stay the same.					
		Describe ho	w the	rate of reaction o	liffers	when the tempera	ature i	s 10°C.	
									[1]
(c)		drochloric aci		ets with iron. uation for this rea	action				
	hy	drochloric acid	+	iron	$\Bigg]  \rightarrow $		+		
(d)				alysts in many ch he term catalyst.	emica	ll reactions.			[2]
									[2]
								[	Total: 9]

7 (a) Fig. 7.1 shows the displayed formula of compound S.

Fig. 7.1

- (i) On Fig. 7.1, draw a circle around the carboxylic acid functional group. [1]
- (ii) Deduce the molecular formula of compound S.

- **(b)** Compound **S** can be converted to acrylic acid. The molecular formula of acrylic acid is C<sub>3</sub>H<sub>4</sub>O<sub>2</sub>.
  - (i) Complete Table 7.1 to calculate the relative molecular mass of acrylic acid.

Table 7.1

atom	number of atoms	relative atomic mass	
carbon	3	12	3 × 12 = 36
hydrogen		1	
oxygen		16	

relative	molecu	lar mas	ss =	 [2	1

[2]

(ii) Acrylic acid is an unsaturated compound.

Describe a test for an unsaturated compound.

test .....

observations .....

	(iii)	When left in the a	ir, acrylic acid forms a	a polymer.		
		State the meaning	g of the term polymer.			
						[2]
	(iv)	Poly(ethene) is al	so a polymer.			
		Choose from the	list the type of polyme	erisation that occurs	when poly(ethene) is	made.
		Draw a circle arou	und your chosen ansv	wer.		
		substitution	n oxidation	neutralisation	addition	[1]
(c)		anoic acid is a carl	•	n of ethanoic acid w	ith sodium hydroxide.	
	(	ethanoic acid +	sodium hydroxide →		+	
						[2]
(d)	Eth	anoic acid can be	converted to ethanol.			
	Nar	me the <b>two</b> produc	ts formed when ethar	nol undergoes comp	lete combustion.	
				and		[2]
					רַ]	Гotal: 13]

Lith	ium bromide is a compound with ionic bonding.	
(a)	State the meaning of the term ionic bond.	
		[2]
(b)	Complete Fig. 8.1 to show:	
	<ul> <li>the electronic configuration of a lithium ion</li> <li>the charge on the ion.</li> </ul>	
	Li	
	Fig. 8.1	[2]
(c)	Deduce the number of protons and neutrons in the bromide ion shown.	
	<sup>79</sup> <sub>35</sub> Br <sup>-</sup>	
	number of protons	
	number of neutrons	 [2]
(d)	Molten lithium bromide is electrolysed using graphite electrodes.	
	State the names of the product at each electrode and give the observations at the positive electrode.	ve
	product at the negative electrode	
	product at the positive electrode	
	observations at the positive electrode	
		 [3]

(e) Fig. 8.2 shows the structure of graphite.




Fig. 8.2

(i)	State the type of bonding in graphite.	
		[1]
(ii)	Explain by referring to Fig. 8.2 why graphite is used as a lubricant.	
		[1]
(iii)	Graphite and diamond are both forms of carbon.	
	State <b>one</b> use of diamond.	
		[1]
	[Total:	12]

## **BLANK PAGE**

### **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

The Periodic Table of Elements

>		₅ He 5	helium 4	10	Ne	neon 20	18	Αr	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	R	radon	118	Og	oganesson -
	:			6	Щ	fluorine 19	17	Cl	chlorine 35.5	35	Ŗ	bromine 80	53	Н	iodine 127	85	¥	astatine -	117	<u>S</u>	tennessine -
>	:			8	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	<u>e</u>	tellurium 128	84	Ъ	moloud –	116	_	livermorium –
>	•			7	Z	nitrogen 14	15	۵	phosphorus 31	33	As	arsenic 75	51	Sb	antimony 122	83	<u>B</u>	bismuth 209	115	Mc	moscovium -
≥	:			9	O	carbon 12	14	S	silicon 28	32	Ge	germanium 73	20	Sn	tin 119	82	Pb	lead 207	114	ŀΙ	flerovium -
	•			2	М	boron 11	13	Αſ	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	lΤ	thallium 204	113	R	mihonium –
							•			30	Zu	zinc 65	48	g	cadmium 112	80	Нg	mercury 201	112	ű	copernicium -
										29	Cn	copper 64	47	Ag	silver 108	62	Αn	gold 197	111	Rg	roentgenium -
Group										28	z	nickel 59	46	Pd	palladium 106	78	చ	platinum 195	110	Ds	darmstadtium -
5										27	ဝိ	cobalt 59	45	뫈	rhodium 103	77	占	iridium 192	109	Ħ	meitnerium -
		- エ	hydrogen 1									iron 56		Ru	ruthenium 101	9/	Os	osmium 190	108	Hs	hassium -
										25	Mn	manganese 55	43	ပ	technetium -	75	Re	rhenium 186	107	Bh	bohrium –
					pol	ass						chromium 52		Mo	molybdenum 96	74	≯	tungsten 184	106	Sg	seaborgium -
			Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	q	niobium 93	73	Б	tantalum 181	105	g O	dubnium -
					ato	rela				22	j	titanium 48	40	Zr	zirconium 91	72	Ξ	hafnium 178	104	Ÿ	rutherfordium -
							•			21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
=	:			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	38	Š	strontium 88	56	Ва	barium 137	88	Ra	radium
_	-			က	:=	lithium 7	11	Na	sodium 23	19	¥	potassium 39	37	&	rubidium 85	55	S	caesium 133	87	ቷ	francium -

71	Γn	lutetium 175	103	۲	lawrencium -
70	Υp	ytterbium 173	102	8	nobelium
69	Tm	thulium 169	101	Md	mendelevium -
89	ш	erbium 167	100	Fm	fermium -
29	웃	holmium 165	66	Es	einsteinium -
99	۵	dysprosium 163	86	ŭ	californium -
65	Д	terbium 159	97	BK	berkelium -
64	Gd	gadolinium 157	96	Cm	curium
63	En	europium 152	92	Am	americium -
62	Sm	samarium 150	94	Pn	plutonium
61	Pm	promethium	93	ď	neptunium -
09	PΝ	neodymium 144	92	$\supset$	uranium 238
69	P	praseodymium 141	91	Ра	protactinium 231
28	Oe	cerium 140	06	드	thorium 232
22	Гa	lanthanum 139	68	Ac	actinium _

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).